

New Relativistic Equations of Motion and Their Comparison with the Equations from the General Relativity

Kostadin Trenčevski*
Faculty of Natural Sciences and Mathematics,
P.O.Box 162, 1000 Skopje, Macedonia

PACS numbers: 04.50.+h, 04.25.-g, 04.80.Cc, 98.80.Jk

Abstract

In this paper is presented the infinitesimally locally Lorentzian theory of gravitation (ILLTG), which predicts the same experimental results which confirm the GR. Motion in a neighborhood of the center of gravitation is considered. The geodesics between the ILLTG and GR are compared. ILLTG differs from GR mainly in the equations of motion and for $v \ll c$ the trajectories are almost the same in both theories. Two tests in the solar system are proposed, such that ILLTG and GR predict different results. The mass and energy with respect to ILLTG is considered. The periastron shift of the binary stars is calculated and a cosmological problem is considered.

1 Introduction

In the paper [8] was presented a new model of gravitational theory, which yields to the same experimental results for motion of a particle in a gravitational field predicted by the general relativity (GR) and which are experimentally verified. In this paper that theory is studied now in many details and a lot new results are presented. In this section will be repeated the basic concepts of the theory given in [8] with some improvements. We convenient to use that $x^4 = ict$ for the time coordinate and orthogonal matrix A with complex elements will mean that $AA^T = I$.

First we underline the main difference between the GR and the proposed theory. Let us start from the metric in GR

$$ds^2 = g_{11}dX^2 + g_{22}dY^2 + g_{33}dZ^2 + g_{44}d(ict)^2,$$

where g_{ij} can approximately be written in the form

$$g_{11} = g_{22} = g_{33} = 1 + 2\frac{V}{c^2}, \quad g_{44} = 1 - 2\frac{V}{c^2}, \quad g_{ij} = 0 \quad (i \neq j).$$

Here V is the gravitational potential, for example $V = \frac{GM}{rc^2}$ in case of gravitation of a spherical body. Note that the coefficients g_{ij} may not be determined very precisely like in the GR, because we shall use them for metric proportions, but not for equations of motion. Thus, the expressions of type c^{-4} and less are neglected. Hence, the experiments for the red shift of the spectral lines and the Shapiro time delay, where the Christoffel symbols are not used, are explained in the same way like in the GR.

This paper was done under the auspices of a CNCSIS Grant.

*Electronic address: kostatre@iunona.pmf.ukim.edu.mk

The above coordinates X, Y, Z, T will be called *ordinary coordinates*. In order to consider the equations of motion, we introduce the so called *orthonormal coordinates* x, y, z, t such that

$$\begin{aligned} dx &= \sqrt{\frac{1}{g_{11}}}dX \approx \left(1 - \frac{V}{c^2}\right)dX, & dy &= \sqrt{\frac{1}{g_{22}}}dY \approx \left(1 - \frac{V}{c^2}\right)dY, & dz &= \sqrt{\frac{1}{g_{33}}}dZ \approx \left(1 - \frac{V}{c^2}\right)dZ, \\ dt &= \sqrt{\frac{1}{g_{44}}}dT \approx \left(1 + \frac{V}{c^2}\right)dT. \end{aligned} \quad (1.1)$$

Further we shall deal mainly with orthonormal coordinates, otherwise it will be emphasized that the coordinates are ordinary. According to the ordinary coordinates, the velocity of the light is equal to $c\left(1 - 2\frac{V}{c^2}\right)$, like in GR, while according to the orthonormal coordinates the velocity of the light is universal constant. Indeed, according to the orthonormal coordinates the space-time in an infinitesimally small neighborhood is flat like in the special theory of relativity (STR), and hence the proposed theory will be called infinitesimally locally Lorentzian theory of gravitation (ILLTG). Note that ILLTG, using orthonormal coordinates, does not describes only gravitation, but also inertial forces (see Ref. [9]). The ordinary coordinates X, Y, Z, T from ILLTG are analogous to the usual coordinates x, y, z, t from the GR, while the orthonormal coordinates from ILLTG mean that the observer is far from gravitation, where the metric is like in the STR.

Instead of the metric g_{ij} we can use only a potential μ . For example, this potential caused by a spherical body with mass M , whose center is at the coordinate origin and which rests with respect to the chosen coordinate system is $\mu = 1 + \frac{GM}{rc^2}$. Thus, the equations (1.1) will be replaced by

$$dx = \mu^{-1}dX, \quad dy = \mu^{-1}dY, \quad dz = \mu^{-1}dZ, \quad dt = \mu dT. \quad (1.1')$$

Further we shall present the basic results about the equations of motion in orthonormal coordinates, and naturally then we should return back in ordinary coordinates as natural. This transformation in an infinitesimally small neighborhood is always possible, but in large we can deal with the nonholonomic coordinates.

If two gravitational fields are given by the gravitational potentials μ_1 and μ_2 , then the superposition of those two fields is a gravitational field with potential $\mu_1\mu_2$. In special case if $\mu_2 = C$ is a constant in the space-time, then the motions of particles should be the same like for the field given by μ_1 .

ILLTG is built using a 4-vector potential U_i . If the magnitude of this vector changes but its direction is a constant, then there exists a gravitational field and the source of gravitational field is moving with 4-vector of velocity $U_i/\|U\|$; if the direction of this vector changes but its magnitude is a constant, then there exist some inertial forces, for example centrifugal. This 4-vector field can be written in the form

$$(U_1, U_2, U_3, U_4) = \frac{\mu}{\sqrt{1 - u^2/c^2}} \left(\frac{u_1}{ic}, \frac{u_2}{ic}, \frac{u_3}{ic}, 1 \right), \quad (1.2)$$

where $\mathbf{u} = (u_1, u_2, u_3)$ is a 3-vector of velocity and μ is the gravitational potential.

Example 1. The STR is characterized by $\mu = \text{const.}$ and $\mathbf{u}$ is a constant vector field.

Example 2. The gravitational field of a spherical body with mass M which is at the origin of the coordinate system is given by the vector field $(0, 0, 0, \mu)$, where $\mu = 1 + \frac{GM}{rc^2}$.

Example 3. If we measure with respect to a noninertial system which rotates around the x^3 axis with angular velocity w , then the 4-vector potential obtains by putting $\mu = 1$ and $\mathbf{u} = (x^2\omega, -x^1\omega, 0)$ [9].

Further we introduce a non-linear connection induced by the 4-vector potential U_i . Let us denote by $S_j^i ds$ the form of connection in the direction of the unit vector $V^i = dx^i/ds$. The necessary and sufficient condition for the metric tensor $g_{ij} = \delta_{ij}$ to be parallel, is

$$S_j^i = -S_i^j. \quad (1.3)$$

Now let us introduce the following tensor

$$F_{ij} = \frac{1}{\mu}(\partial U_i / \partial x^j - \partial U_j / \partial x^i). \quad (1.4)$$

If we denote by V_i the 4-vector of velocity of the moving particle, i.e.

$$(V_i) = \left(\frac{v_1}{ic\sqrt{1-v^2/c^2}}, \frac{v_2}{ic\sqrt{1-v^2/c^2}}, \frac{v_3}{ic\sqrt{1-v^2/c^2}}, \frac{1}{\sqrt{1-v^2/c^2}} \right), \quad (1.5)$$

one can easily verify that the following simple equations

$$dV_i/ds = F_{ij}V_j, \quad (1.6)$$

which are analogous to the Lorentz force in the theory of electromagnetism, contain inertial accelerations (Coriolis acceleration, centrifugal and transversely acceleration) and gravitation acceleration for Newton approach. In order to obtain a precise theory, we consider the following two tensors

$$\phi_{ij} = \frac{1}{2}F_{ij} - \frac{1}{2}\mu^{-2}(U_i U_k F_{jk} - U_j U_k F_{ik}), \quad (1.7)$$

$$P_{ij} = \delta_{ij} - \frac{1}{\mu + U_s V_s}(\mu V_i V_j + V_i U_j + U_i V_j + \frac{1}{\mu} U_i U_j) + \frac{2}{\mu} U_j V_i. \quad (1.8)$$

These two tensors together with F_{ij} are invariant under the gauge transformation $\mu \rightarrow C\mu$, where C is a constant. The tensors F_{ij} and ϕ_{ij} are antisymmetric and P_{ij} is an orthogonal matrix. It can be verified by using the identities $U_i U_i = \mu^2$ and $V_i V_i = 1$. Now the required connection S_{ij} is introduced by

$$S_{ij} = -P_{ri} \phi_{rk} P_{kj}, \quad (1.9)$$

or in matrix form $S = -P^t \phi P$. Although this connection seems to be very complicated, now we will show the opposite. Indeed, assume that $(U_i) = (0, 0, 0, 1)$ at the considered point. Then

$$F = \begin{bmatrix} 0 & -2i\omega_3/c & 2i\omega_2/c & -a_1/c^2 \\ 2i\omega_3/c & 0 & -2i\omega_1/c & -a_2/c^2 \\ -2i\omega_2/c & 2i\omega_1/c & 0 & -a_3/c^2 \\ a_1/c^2 & a_2/c^2 & a_3/c^2 & 0 \end{bmatrix}, \quad (1.10)$$

$$\phi = \begin{bmatrix} 0 & -i\omega_3/c & i\omega_2/c & -a_1/c^2 \\ i\omega_3/c & 0 & -i\omega_1/c & -a_2/c^2 \\ -i\omega_2/c & i\omega_1/c & 0 & -a_3/c^2 \\ a_1/c^2 & a_2/c^2 & a_3/c^2 & 0 \end{bmatrix}, \quad (1.11)$$

where

$$\mathbf{a} = (a_1, a_2, a_3) = c^2 \text{grad } U_4 + \partial \mathbf{u} / \partial t \quad \text{and} \quad \mathbf{w} = -\frac{1}{2} \text{rot } \mathbf{u} \quad (1.12)$$

represent the 3-vector of acceleration and the 3-vector of angular velocity. In that special case $(U_i) = (0, 0, 0, 1)$, the tensor P is given by

$$P = \begin{bmatrix} 1 - \frac{1}{\nu} V_1^2 & -\frac{1}{\nu} V_1 V_2 & -\frac{1}{\nu} V_1 V_3 & V_1 \\ -\frac{1}{\nu} V_2 V_1 & 1 - \frac{1}{\nu} V_2^2 & -\frac{1}{\nu} V_2 V_3 & V_2 \\ -\frac{1}{\nu} V_3 V_1 & -\frac{1}{\nu} V_3 V_2 & 1 - \frac{1}{\nu} V_3^2 & V_3 \\ -V_1 & -V_2 & -V_3 & V_4 \end{bmatrix}, \quad (1.13)$$

where V_1, V_2, V_3, V_4 are given by (1.5) and $\nu = 1 + V_4$. Note that the matrix P represents just a Lorentz transformation for the 3-vector $-\mathbf{v} = (-v_1, -v_2, -v_3)$. Now the connection S_{ij} becomes much more clear and acceptable.

Remark. Although ϕ is a tensor, its components have physical meaning given by (1.11) only for $U_1 = U_2 = U_3 = 0$. On the other hand, the tensor F defined by (1.4) always has components given approximately by (1.10). In case of gravitational field, $U_1 = U_2 = U_3 = 0$ means that the source of gravitation is in rest with respect to the chosen coordinate system. In this system it is $F_{ij} = \phi_{ij}$ assuming that $\mathbf{w} = 0$, and hence it is true in each coordinate system.

Note that the introduced connection S_{ij} is non-linear. The first step of approximation yields to the following linear connection $\Gamma_k = (\Gamma_{jk}^i)$ such that $S_j^i \approx \Gamma_{js}^i V_s$, where

$$\Gamma_1 = \begin{bmatrix} 0 & a_2/c^2 & a_3/c^2 & 0 \\ -a_2/c^2 & 0 & 0 & -iw_3/c \\ -a_3/c^2 & 0 & 0 & iw_2/c \\ 0 & iw_3/c & -iw_2/c & 0 \end{bmatrix}, \quad (1.14a)$$

$$\Gamma_2 = \begin{bmatrix} 0 & -a_1/c^2 & 0 & iw_3/c \\ a_1/c^2 & 0 & a_3/c^2 & 0 \\ 0 & -a_3/c^2 & 0 & -iw_1/c \\ -iw_3/c & 0 & iw_1/c & 0 \end{bmatrix}, \quad (1.14b)$$

$$\Gamma_3 = \begin{bmatrix} 0 & 0 & -a_1/c^2 & -iw_2/c \\ 0 & 0 & -a_2/c^2 & iw_1/c \\ a_1/c^2 & a_2/c^2 & 0 & 0 \\ iw_2/c & -iw_1/c & 0 & 0 \end{bmatrix}, \quad (1.14c)$$

$$\Gamma_4 = \begin{bmatrix} 0 & iw_3/c & -iw_2/c & a_1/c^2 \\ -iw_3/c & 0 & iw_1/c & a_2/c^2 \\ iw_2/c & -iw_1/c & 0 & a_3/c^2 \\ -a_1/c^2 & -a_2/c^2 & -a_3/c^2 & 0 \end{bmatrix}, \quad (1.14d)$$

and where we assumed that $(U_i) = (0, 0, 0, 1)$ at the considered point. The Newton's approximation obtains by replacing the components Γ_{jk}^i for $i, j, k \in \{1, 2, 3\}$ with zero. Using this linear connection the curvature tensor is easy to be calculated and in [8] it is shown that the Einstein equations of gravitational field

$$R_{ij} - \frac{1}{2}g_{ij}R = 8G\pi c^{-2}T_{ij} \quad (1.15)$$

are satisfied, where the energy-momentum tensor T_{ij} is given by $T_{ij} = \rho U_i U_j$. It is also shown in [8] that the following equations

$$(\mu F_{ij})_{;k} + (\mu F_{jk})_{;i} + (\mu F_{ki})_{;j} = 0, \quad (1.16)$$

$$(\mu F_{ij})_{;j} = 4G\pi c^{-2}J_i, \quad (1.17)$$

analogous to the Maxwell equations, also hold at that order of approximation. Here $J_i = \rho U_i$ is the 4-current density for gravitation, analogous to the 4-current density for the electromagnetism.

2 Some results for the geodesics according to the ILLTG

In this section the planetary orbits in the solar system as well as the trajectories of the light rays will be considered. Our coordinate origin will be chosen to be at the center of the Sun. The source of gravitation is in rest to our coordinates system, and we can put $U_1 = U_2 = U_3 = 0$. We shall assume that the mass of the moving body is negligible with respect to the Sun. The general equations of motion of a body with nonzero mass are given in section 6. We shall deal with the orthonormal coordinates.

Let $\mu = 1 + \frac{GM}{rc^2}$, where M is the mass of the Sun. Then

$$(F_{ij}) = \begin{bmatrix} 0 & 0 & 0 & \frac{GM}{r^3 c^2} x \\ 0 & 0 & 0 & \frac{GM}{r^3 c^2} y \\ 0 & 0 & 0 & \frac{GM}{r^3 c^2} z \\ -\frac{GM}{r^3 c^2} x & -\frac{GM}{r^3 c^2} y & -\frac{GM}{r^3 c^2} z & 0 \end{bmatrix} \quad (2.1)$$

and the equations of motion $\frac{dV_i}{ds} = -P_{ri}\phi_{rk}P_{kj}$ for $i = 1, 2, 3, 4$ without approximation reduce to [8]

$$\frac{d^2 x}{dt^2} = \frac{GM}{\mu r^3} \left[(\beta^{-2} - 2)x + \frac{v_x}{c^2} (xv_x + yv_y) \left(2 + \frac{1}{\beta(\beta + 1)} \right) \right], \quad (2.2a)$$

$$\frac{d^2 y}{dt^2} = \frac{GM}{\mu r^3} \left[(\beta^{-2} - 2)y + \frac{v_y}{c^2} (xv_x + yv_y) \left(2 + \frac{1}{\beta(\beta + 1)} \right) \right], \quad (2.2b)$$

$$\beta - \ln \mu = C_1 (= \text{const.}), \quad (2.2c)$$

where $\beta = \left(1 - \frac{v^2}{c^2} \right)^{-1/2}$. Without loss of generality we assumed that $v_z = 0$ and that the particle moves in the xy -plane. Thus we omit the equation for $i = 3$.

Remark. Using the formulas (1.1'), and the fact that μ is a scalar which does not depend from the coordinate system (orthonormal or ordinary), it is easy to see that in ordinary coordinates the equations of motion (2.2) become:

$$\frac{d^2 X}{dT^2} = \frac{2}{\mu} \frac{d\mu}{dT} \frac{dX}{dT} + \mu^3 \frac{d^2 x}{dt^2}, \quad (2.2'a)$$

$$\frac{d^2 Y}{dT^2} = \frac{2}{\mu} \frac{d\mu}{dT} \frac{dY}{dT} + \mu^3 \frac{d^2 y}{dt^2} \quad (2.2'b)$$

$$\beta - \ln \mu = C_1 (= \text{const.}), \quad (2.2'c)$$

where $\frac{d^2 x}{dt^2}$ and $\frac{d^2 y}{dt^2}$ are given by (2.2a,b). Note that the equation (2.2c) remains unchanged. Indeed, μ appears to be the same in both coordinate systems and also the quotient v^2/c^2 takes the same value in both coordinate systems.

Remark. Note that if we have not only a gravitational field with a potential μ , but we also have a 3-vector of acceleration (a_x, a_y, a_z) and a 3-vector of angular velocity (w_x, w_y, w_z) , then in orthonormal coordinates it is

$$\frac{dv_x}{dt} = \frac{1}{\mu} \left[(2 - \beta^{-2})a_x - \frac{v_x}{c^2} (a_x v_x + a_y v_y + a_z v_z) \left(2 + \frac{1}{\beta(\beta + 1)} \right) + 2(v_y w_z - v_z w_y) \right], \quad (2.3a)$$

$$\frac{dv_y}{dt} = \frac{1}{\mu} \left[(2 - \beta^{-2})a_y - \frac{v_y}{c^2} (a_x v_x + a_y v_y + a_z v_z) \left(2 + \frac{1}{\beta(\beta + 1)} \right) + 2(v_z w_x - v_x w_z) \right], \quad (2.3b)$$

$$\frac{dv_z}{dt} = \frac{1}{\mu} \left[(2 - \beta^{-2})a_z - \frac{v_z}{c^2} (a_x v_x + a_y v_y + a_z v_z) \left(2 + \frac{1}{\beta(\beta + 1)} \right) + 2(v_x w_y - v_y w_x) \right], \quad (2.3c)$$

$$\frac{d}{dt} \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{1}{\mu c^2} (a_x v_x + a_y v_y + a_z v_z). \quad (2.3d)$$

The acceleration which is caused by the gravitational potential μ is included in the 3-vector of acceleration (a_x, a_y, a_z) .

Now, using that

$$\frac{d\varphi}{dt} = \frac{d}{dt} \arctan \frac{y}{x} = \frac{v_y x - v_x y}{r^2},$$

from (2.2a) and (2.2b), we obtain

$$\begin{aligned} \frac{d}{dt} \left(r^2 \frac{d\varphi}{dt} \right) &= \frac{d^2 y}{dt^2} x - \frac{d^2 x}{dt^2} y = \frac{GM}{\mu r^3 c^2} (v_y x - v_x y) (x v_x + y v_y) \left(2 + \frac{1}{\beta(\beta+1)} \right) = \\ &= -r^2 \frac{d\varphi}{dt} \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) \frac{d}{dt} \left(\frac{1}{r} \right). \end{aligned}$$

Two cases will be considered.

(i) Assume that $v \ll c$, i.e. consider a planetary orbit. Then $2 + \frac{1}{\beta(\beta+1)} \approx 2,5$ and hence

$$\frac{d}{dt} \left(r^2 \frac{d\varphi}{dt} \right) = -\frac{5}{2} \frac{GM}{c^2} \left(r^2 \frac{d\varphi}{dt} \right) \frac{d}{dt} \left(\frac{1}{r} \right),$$

by neglecting the expressions of order c^{-4} . The solution of the previous equation is

$$r^2 \frac{d\varphi}{dt} = C_2 \exp \left(\frac{-5}{2} \frac{GM}{rc^2} \right), \quad C_2 = \text{const.} \quad (2.4)$$

Further, using the metric $(dr)^2 + r^2(d\varphi)^2 - c^2 t^2 = ds^2$ in orthonormal coordinates, we obtain

$$\left(\frac{dr}{dt} \right)^2 + \left(r \frac{d\varphi}{dt} \right)^2 = v^2, \quad \left(r^{-2} \frac{dr}{d\varphi} \right)^2 + r^{-2} = v^2 \left(r^2 \frac{d\varphi}{dt} \right)^{-2},$$

and get

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = v^2 C_2^{-2} \exp \left(\frac{5GM\rho}{c^2} \right), \quad (2.5)$$

where $\rho = r^{-1}$. From (2.2c) we obtain approximately

$$\left(1 - \frac{v^2}{c^2} \right)^{-1/2} - \frac{GM}{rc^2} \left(1 - \frac{1}{2} \frac{GM}{rc^2} \right) = 1 + c^{-2} C_3,$$

where $C_3 = c^2(C_1 - 1) = \text{const.}$ Hence we obtain

$$\begin{aligned} v^2 &= c^2 \left[1 - \left(1 + c^{-2} \left(C_3 + GM\rho \left(1 - \frac{1}{2} \frac{GM\rho}{c^2} \right) \right) \right)^{-2} \right] = \\ &= 2C_3 + 2GM\rho \left(1 - \frac{1}{2} \frac{GM\rho}{c^2} \right) - 3c^{-2} C_3^2 - 6C_3 \frac{GM\rho}{c^2} - 3 \frac{G^2 M^2 \rho^2}{c^2}. \end{aligned}$$

Replacing this value of v^2 into (2.5), after neglecting the summands of order c^{-4} , we obtain the following differential equation

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = A + B\rho + \frac{6G^2 M^2}{c^2 C_2^2} \rho^2 \quad (2.6)$$

and hence

$$d\varphi = - \left(1 + \frac{3G^2 M^2}{c^2 C_2^2} \right) \frac{d\rho}{\sqrt{-\rho^2 + p\rho - q}},$$

where p and q are constants. Indeed, $p = \rho_1 + \rho_2$, $q = \rho_1 \rho_2$, where $\rho_1 = \frac{1}{r_1}$, $\rho_2 = \frac{1}{r_2}$ and r_1 and r_2 are extreme distances ($r_1 < r_2$) from the planet to the center of the Sun. It is well known that $C_2 = \sqrt{GMa(1-\epsilon^2)}$, where $a = (r_1 + r_2)/2$ and ϵ is the eccentricity of the approximate ellipse. Thus,

$$d\varphi = - \left(1 + \frac{3GM}{ac^2(1-\epsilon^2)} \right) \frac{d\rho}{\sqrt{-\rho^2 + p\rho - q}}. \quad (2.7)$$

Hence we obtain

$$\varphi = -\left(1 + \frac{3GM}{ac^2(1-\epsilon^2)}\right) \left(\arcsin \frac{\rho - \frac{p}{2}}{\sqrt{\frac{p^2}{4} - q}} - C_4\right) = -\left(1 + \frac{3GM}{ac^2(1-\epsilon^2)}\right) \left(\arcsin \frac{2\rho - \rho_1 - \rho_2}{\rho_1 - \rho_2} - C_4\right).$$

Assume that $\varphi = 0$ if $\rho = \rho_1$, and hence $C_4 = \pi/2$. This equation can be solved by ρ and its solution is

$$\rho = \frac{\rho_1 + \rho_2}{2} + \frac{\rho_1 - \rho_2}{2} \cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1-\epsilon^2)} \right) \right]. \quad (2.8)$$

Now it follows that the angle between the points for which $\rho = \rho_1$ and $\rho = \rho_2$ is equal to $\pi + \frac{3GM\pi}{ac^2(1-\epsilon^2)}$ and hence the angle for the perihelion shift is equal to $\Delta\varphi = \frac{6GM\pi}{ac^2(1-\epsilon^2)}$.

(ii) Let us consider the trajectory of a light ray near the Sun. We denote by R the radius of the Sun. In this case $\beta = \infty$,

$$\frac{d}{dt} \left(r^2 \frac{d\varphi}{dt} \right) = -2 \frac{GM}{c^2} \left(r^2 \frac{d\varphi}{dt} \right) \frac{d}{dt} \left(\frac{1}{r} \right)$$

and its solution is

$$r^2 \frac{d\varphi}{dt} = C_2 \exp \left(-2 \frac{GM}{rc^2} \right), \quad C_2 = \text{const}. \quad (2.9)$$

Analogously to (2.5) we obtain

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = v^2 C_2^{-2} \exp \left(\frac{4GM\rho}{c^2} \right)$$

and replacing $v = c$, we get

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = c^2 C_2^{-2} \exp \left(\frac{4GM\rho}{c^2} \right). \quad (2.10)$$

We shall prove now that the light has a constant velocity c in a gravitational field, using orthonormal coordinates. Summing up the equation (2.2a) multiplied by $2v_x$ and (2.2b) multiplied by $2v_y$ we obtain

$$\begin{aligned} \frac{dv^2}{dt} &= \frac{2GM}{\mu r^3} \left[-\left(1 + \frac{v^2}{c^2}\right) (xv_x + yv_y) + (xv_x + yv_y) \left(2 + \frac{1}{\beta(\beta+1)}\right) \frac{v^2}{c^2} \right] = \\ &= -\frac{2GM}{\mu r^3} (xv_x + yv_y) \frac{\left(1 - \frac{v^2}{c^2}\right)^{3/2}}{1 + \sqrt{1 - \frac{v^2}{c^2}}}. \end{aligned}$$

Hence, if $v^2 = c^2$ at one point, then $v^2 = c^2$ holds on the whole trajectory, because $v^2 = c^2$ satisfies the previous differential equation and the initial condition.

If $r = R$, then $R \frac{d\varphi}{dt} = c$ and from (2.9) we get $C_2 = Rc \exp \left(\frac{2GM}{Rc^2} \right)$. By replacing this value of C_2 into (2.10) we get

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = \frac{1}{R^2} \exp \left(\frac{4GM}{c^2} \left(\rho - \frac{1}{R} \right) \right). \quad (2.11)$$

By neglecting the expressions of order c^{-4} we obtain

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = \frac{1}{R^2} - \frac{4GM}{R^3 c^2} + \frac{4GM}{R^2 c^2} \rho, \quad (2.12)$$

which is equation of a hyperbola. Note that according to the Newton's theory the trajectory is a different hyperbola: $\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = \frac{1}{R^2} - \frac{2GM}{R^3 c^2} + \frac{2GM}{R^2 c^2} \rho$.

From (2.12) φ can be determined as a function of ρ :

$$\varphi = \arccos \frac{\rho R^2 c^2 - 2GM}{Rc^2 - 2GM}, \quad (2.13)$$

such that $\varphi = 0$ if $\rho = \frac{1}{R}$. Hence from (2.13) we obtain

$$\rho = \frac{2GM}{R^2 c^2} + \left(\frac{1}{R} - \frac{2GM}{R^2 c^2} \right) \cos \varphi. \quad (2.14)$$

It is easy to conclude from (2.13) that the angle of deflection of a light ray near the Sun is equal to $\frac{4GM}{Rc^2}$. This conclusion can be obtained also by the following way. Assume that the light ray is parallel to the x -axis at $(0, R, 0, 0)$. Let us put $\frac{dx}{dt} = v \cos \alpha$, $\frac{dy}{dt} = -v \sin \alpha$, $x = vt$, $y = R$ and $r = \sqrt{R^2 + v^2 t^2}$ in (2.2a) and (2.2b). Then

$$\frac{dv}{dt} \cos \alpha - v \sin \alpha \frac{d\alpha}{dt} = \frac{GM}{\mu r^3} \left[-\left(1 + \frac{v^2}{c^2}\right) vt + \frac{v \cos \alpha}{c^2} (xv_x + yv_y) \left(2 + \frac{1}{\beta(\beta+1)}\right) \right], \quad (2.15a)$$

$$-\frac{dv}{dt} \sin \alpha - v \cos \alpha \frac{d\alpha}{dt} = \frac{GM}{\mu r^3} \left[-\left(1 + \frac{v^2}{c^2}\right) R - \frac{v \sin \alpha}{c^2} (xv_x + yv_y) \left(2 + \frac{1}{\beta(\beta+1)}\right) \right]. \quad (2.15b)$$

Summing up the first equation multiplied by $\sin \alpha$ and the second equation multiplied by $\cos \alpha$, we obtain

$$v \frac{d\alpha}{dt} = \frac{GM}{\mu r^3} \left(1 + \frac{v^2}{c^2}\right) (R \cos \alpha + vt \sin \alpha).$$

By putting $v = c$, $\mu \approx 1$ and $R \cos \alpha + vt \sin \alpha \approx R$, we get

$$\begin{aligned} \frac{d\alpha}{dt} &= \frac{2GMR}{c(R^2 + c^2 t^2)^{3/2}}, \\ \Delta\alpha &= \int_{-\infty}^{\infty} \frac{2GMR dt}{c(R^2 + c^2 t^2)^{3/2}} = \frac{4GM}{Rc^2}. \end{aligned} \quad (2.16)$$

Although we deduced the main formulas $\Delta\varphi = \frac{6GM\pi}{ac^2(1-\epsilon^2)}$ and $\Delta\alpha = \frac{4GM}{Rc^2}$ by using orthonormal coordinates, they remain unchanged if we use ordinary coordinates.

To the end of this section we shall consider the motion of a particle (with zero mass) under the gravitation of k gravitational objects. Indeed, we have in special case a motion in the solar system with perturbations of the planets and other gravitational bodies. Let m_i be the mass of the i -th gravitational object and r_i be the distance from its center to the particle. Further, let the 3-vector of velocity of the particle is $v = (v_x, v_y, v_z)$ and the 3-vector of velocity of the i -th gravitational object is $u_i = (u_{ix}, u_{iy}, u_{iz})$ and let us put

$$U_i = -c^2 \ln \left(1 + \frac{Gm_i}{r_i c^2} \right). \quad (2.17)$$

Let us denote the 4-vector of velocity of the i -th object by

$$(U_{i1}, U_{i2}, U_{i3}, U_{i4}) = \left(\frac{\frac{u_{ix}}{ic}}{\sqrt{1 - \frac{u_i^2}{c^2}}}, \frac{\frac{u_{iy}}{ic}}{\sqrt{1 - \frac{u_i^2}{c^2}}}, \frac{\frac{u_{iz}}{ic}}{\sqrt{1 - \frac{u_i^2}{c^2}}}, \frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}} \right),$$

such that the 4-vector potential is $\left(1 + \frac{Gm_i}{r_i c^2}\right) (U_{i1}, U_{i2}, U_{i3}, U_{i4})$.

The tensors F_{pq} and ϕ_{pq} considered for the i -th object are

$$F_{pq}^i = \phi_{pq}^i = \frac{1}{c^2} \left[\frac{\partial U_i}{\partial x_p} U_{iq} - \frac{\partial U_i}{\partial x_q} U_{ip} \right]$$

(do not sum over i). Now, analogously to (2.3) the equations of motion under the gravitation of the i -th object is given by

$$\frac{dv_x}{dt} = \left[(2 - \beta^{-2}) a_x - \frac{v_x}{c^2} (a_x v_x + a_y v_y + a_z v_z) \left(2 + \frac{1}{\beta(\beta+1)}\right) + 2(v_y w_z - v_z w_y) \right], \quad (2.18a)$$

$$\frac{dv_y}{dt} = \left[(2 - \beta^{-2})a_y - \frac{v_y}{c^2}(a_x v_x + a_y v_y + a_z v_z)(2 + \frac{1}{\beta(\beta+1)}) + 2(v_z w_x - v_x w_z) \right], \quad (2.18b)$$

$$\frac{dv_z}{dt} = \left[(2 - \beta^{-2})a_z - \frac{v_z}{c^2}(a_x v_x + a_y v_y + a_z v_z)(2 + \frac{1}{\beta(\beta+1)}) + 2(v_x w_y - v_y w_x) \right], \quad (2.18c)$$

$$\frac{d}{dt} \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{1}{c^2}(a_x v_x + a_y v_y + a_z v_z), \quad (2.18d)$$

where

$$a_x = \left[U_{i1} \frac{\partial U_i}{\partial x_4} - U_{i4} \frac{\partial U_i}{\partial x_1} \right], \quad a_y = \left[U_{i2} \frac{\partial U_i}{\partial x_4} - U_{i4} \frac{\partial U_i}{\partial x_2} \right], \quad a_z = \left[U_{i3} \frac{\partial U_i}{\partial x_4} - U_{i4} \frac{\partial U_i}{\partial x_3} \right],$$

$$w_x = \frac{i}{c} \left[U_{i3} \frac{\partial U_i}{\partial x_2} - U_{i2} \frac{\partial U_i}{\partial x_3} \right], \quad w_y = \frac{i}{c} \left[U_{i1} \frac{\partial U_i}{\partial x_3} - U_{i3} \frac{\partial U_i}{\partial x_1} \right], \quad w_z = \frac{i}{c} \left[U_{i2} \frac{\partial U_i}{\partial x_1} - U_{i1} \frac{\partial U_i}{\partial x_2} \right].$$

If the potential U_i is not time dependent, then

$$a_x = -\frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}} \frac{\partial U_i}{\partial x}, \quad a_y = -\frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}} \frac{\partial U_i}{\partial y}, \quad a_z = -\frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}} \frac{\partial U_i}{\partial z}.$$

Further, after some algebraic transformations, the final equations for the motion of the particle under the gravitation of all objects become

$$\begin{aligned} \frac{dv_x}{dt} = & \sum_{i=1}^k \left[\frac{\partial U_i}{\partial x} (\beta^{-2} - 2 + 2 \frac{u_{ix} v_x + u_{iy} v_y + u_{iz} v_z}{c^2}) + \frac{\partial U_i}{\partial x} \frac{M v_x^2 - 2 u_{ix} v_x}{c^2} + \right. \\ & \left. + \frac{\partial U_i}{\partial y} \frac{M v_x v_y - 2 u_{ix} v_y}{c^2} + \frac{\partial U_i}{\partial z} \frac{M v_x v_z - 2 u_{ix} v_z}{c^2} \right] \frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}}, \end{aligned} \quad (2.19a)$$

$$\begin{aligned} \frac{dv_y}{dt} = & \sum_{i=1}^k \left[\frac{\partial U_i}{\partial y} (\beta^{-2} - 2 + 2 \frac{u_{ix} v_x + u_{iy} v_y + u_{iz} v_z}{c^2}) + \frac{\partial U_i}{\partial x} \frac{M v_x v_y - 2 u_{iy} v_x}{c^2} + \right. \\ & \left. + \frac{\partial U_i}{\partial y} \frac{M v_y^2 - 2 u_{iy} v_y}{c^2} + \frac{\partial U_i}{\partial z} \frac{M v_y v_z - 2 u_{iy} v_z}{c^2} \right] \frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}}, \end{aligned} \quad (2.19b)$$

$$\begin{aligned} \frac{dv_z}{dt} = & \sum_{i=1}^k \left[\frac{\partial U_i}{\partial z} (\beta^{-2} - 2 + 2 \frac{u_{ix} v_x + u_{iy} v_y + u_{iz} v_z}{c^2}) + \frac{\partial U_i}{\partial x} \frac{M v_x v_z - 2 u_{iz} v_x}{c^2} + \right. \\ & \left. + \frac{\partial U_i}{\partial y} \frac{M v_y v_z - 2 u_{iz} v_y}{c^2} + \frac{\partial U_i}{\partial z} \frac{M v_z^2 - 2 u_{iz} v_z}{c^2} \right] \frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}}, \end{aligned} \quad (2.19c)$$

$$\frac{d}{dt} \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} + \frac{1}{c^2} \sum_{i=1}^k \frac{1}{\sqrt{1 - \frac{u_i^2}{c^2}}} \left(\frac{\partial U_i}{\partial x} v_x + \frac{\partial U_i}{\partial y} v_y + \frac{\partial U_i}{\partial z} v_z \right) = 0, \quad (2.19d)$$

where $M = 2 + \frac{1}{\beta(\beta+1)}$. It is important to emphasize that $\frac{\partial U_i}{\partial x} v_x + \frac{\partial U_i}{\partial y} v_y + \frac{\partial U_i}{\partial z} v_z$ is not a total differential of U_i , because U_i depends also on the coordinates of the i -th body.

3 Representations of the geodesics near the Schwarzschild radii according to the ILLTG

In section 2 we considered the geodesics in polar coordinates with approximation, neglecting the expressions of order c^{-4} and smaller. In this section we shall consider the geodesics in polar coordinates without approximation, using orthonormal coordinates. Starting from the equation (section 2)

$$\frac{d}{dt} \left(r^2 \frac{d\varphi}{dt} \right) = - \left(r^2 \frac{d\varphi}{dt} \right) \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) \frac{d}{dt} \left(\frac{1}{r} \right),$$

we obtain

$$\frac{d}{dt} \left(\ln \left(r^2 \frac{d\varphi}{dt} \right) \right) = - \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) \frac{d}{dt} \left(\frac{1}{r} \right).$$

Replacing in this equation that $\beta = C_1 + \ln \left(1 + \frac{GM}{rc^2} \right) = C_1 + \ln \left(1 + \frac{GM\rho}{c^2} \right)$ as a function of ρ , we obtain

$$\begin{aligned} \frac{d}{dt} \left(\ln \left(r^2 \frac{d\varphi}{dt} \right) \right) &= \frac{d}{dt} \int \left(- \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) \right) d \left(\frac{1}{r} \right), \\ r^2 \frac{d\varphi}{dt} &= C_2 \exp \left(- \int \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) d\rho \right) \end{aligned}$$

and further like in section 2 we obtain

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = v^2 C_2^{-2} \exp \left(\int \frac{2GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) d\rho \right),$$

where

$$v^2 = c^2 \left[1 - \left(C_1 + \ln \mu \right)^{-2} \right], \quad \mu = 1 + \frac{GM\rho}{c^2}.$$

The constant C_2^{-2} will be determined, assuming that $\frac{d\rho}{d\varphi} = 0$ at the considered point. Indeed,

$$\rho_0^2 = v_0^2 C_2^{-2} \exp \left(\int \frac{2GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) d\rho \Big|_{\rho=\rho_0} \right),$$

and hence

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = v^2 \frac{\rho_0^2}{v_0^2} \exp \left(\int 2 \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) d\rho - \int 2 \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) d\rho \Big|_{\rho=\rho_0} \right). \quad (3.1)$$

We convenient to use the short notations

$$\int 2 \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) d\rho = I \quad \text{and} \quad \int 2 \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) d\rho \Big|_{\rho=\rho_0} = I_0.$$

Now, (3.1) can be written in the following forms

$$\frac{\frac{1}{v^2} \left(\frac{d\rho}{d\varphi} \right)^2 + \frac{\rho^2}{v^2}}{e^I} = \frac{\rho_0^2}{v_0^2} \frac{1}{e^{I_0}}, \quad (3.2)$$

$$\left(\frac{d\rho}{d\varphi} \right)^2 = \left[\frac{\rho_0^2}{v_0^2} \frac{1}{e^{I_0}} - \frac{\rho^2}{v^2} \frac{1}{e^I} \right] v^2 e^I. \quad (3.3)$$

Let us denote by f the following *characteristic function*

$$f(\rho) = \frac{\rho^2}{v^2 e^I}. \quad (3.4)$$




Fig. 1

Note that $v^2 = c^2[1 - (C_1 + \ln \mu)^{-2}]$ and we can consider the function (3.4) only for those ρ , for which $v^2 \geq 0$. Two cases are possible and we shall consider them separately.

(i) Assume that $C_1 \geq 1$. The case $C_1 = \infty$, i.e. $v = c$ is included. Physically $C_1 > 1$ means that if the particle left the gravitational field and goes into infinity ($r = \infty$), then it will move with velocity v_0 such that $C_1 = \left(1 - \frac{v_0^2}{c^2}\right)^{-1/2}$. Specially if $C_1 = 1$, then $v_0 = 0$. The characteristic function $f(\rho)$ is defined for each $\rho > 0$ and obviously $\lim_{\rho \rightarrow 0} f(\rho) = 0$. Since $2 \leq 2 + \frac{1}{\beta(\beta+1)} < 2\frac{1}{2}$, we get $\mu^4 \leq e^I < \mu^5$. Thus, if $\rho \rightarrow \infty$, then $v \rightarrow c$ and using the previous inequalities we obtain $\lim_{\rho \rightarrow \infty} f(\rho) = 0$. We shall find the maximal value of $f(\rho)$. Indeed, $f'(\rho) = 0$ can easily be written in an equivalent form

$$\left[\frac{dv}{d\rho} + \frac{vGM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) \right] \rho^2 = v\rho.$$

By differentiating the equality $\beta - \ln \mu = C_1$, it is easy to get $\frac{dv}{d\rho} = \frac{GM}{\mu} \frac{1}{v} \left(1 - \frac{v^2}{c^2} \right)^{3/2}$. Thus, $f'(\rho) = 0$ if and only if

$$\frac{GM\rho}{\mu} \frac{1}{v} \left(1 - \frac{v^2}{c^2} \right)^{3/2} + v \frac{GM\rho}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)} \right) = v$$

and after some algebraic transformations it is equivalent to

$$v^2 = GM\rho. \quad (3.5)$$

The physical meaning of (3.5) is very simple. Indeed, it is easy to verify that the equations (2.2) are verified if a particle moves with velocity $v = \sqrt{GM\rho}$ on a distance $r = \frac{1}{\rho}$ on a circle. It is just like in the Newton's theory, but remember that $C_1 > 1$ which is analogous with $\frac{1}{2}v^2 - GM\rho > 0$ in the Newton's theory and according to it the trajectory is hyperbola, but not a circle. We shall prove that this effect in ILLTG is a "relativistic effect". The equation (3.5) is equivalent to

$$C_1 = \frac{1}{\sqrt{1 - \frac{GM\rho}{c^2}}} - \ln \left(1 + \frac{GM\rho}{c^2} \right). \quad (3.6)$$

The graphic of the right side of (3.6) is given on Fig. 1.

Each of these circles has radius in the interval $\left[\frac{c^2}{GM}, \frac{1}{0,5030} \frac{c^2}{GM} \right]$ and will be called *Schwarzschild radius* which corresponds to the chosen value $C_1 > 1$. We shall denote these radii by index s . The graphic of the function $f(\rho)$ is given on figure 2. On fig. 3 is given the graphic of $f(\rho)$ when $C_1 = 1$. The graphics are compared with the corresponding Newton's case $f_N(\rho) = \frac{\rho^2}{v^2} = \frac{\rho^2}{const. + 2GM\rho}$.




Fig.2




Fig.3

Now according to (3.2) we are able to draw some conclusions for the trajectory. Let be given ρ , v and $\frac{d\rho}{d\varphi}$ at a chosen point. Since ρ and v are known, the constant C_1 (≥ 1) is known and also the function $f(\rho)$ is known. We have the following options and the conclusions follow by Propositions 1 and 2 at the end of this section.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} > f(\rho_s)$, then the equation (3.2) has no solutions for ρ_0 and hence $\frac{d\rho}{d\varphi} = 0$ will never be achieved and the particle will tend to the center $r = 0$. This easily can happen for large initial values of $\frac{d\rho}{d\varphi}$.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} = f(\rho_s)$, then the particle will move on a circle.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} < f(\rho_s)$, and $\rho < \rho_s$, i.e. $r > r_s$, then the trajectory of the particle will be out of the Schwarzschild sphere (S.s.), it will look like a hyperbola and $\frac{d\rho}{d\varphi} = 0$ will be achieved once.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} < f(\rho_s)$, but $\rho > \rho_s$, i.e. $r < r_s$, then the trajectory of the particle will be only inside of the S.s., and it will tend to the center $r = 0$. The condition $\frac{d\rho}{d\varphi} = 0$ will be achieved once.

According to the previous discussion we see that the S.s. with radius $r = r_s$ is a sphere of separation. We shall consider further in more details the motion of a light ray. In this case $C_1 = \infty$, $v = c$ and $\rho_s = \frac{c^2}{GM}$. On fig. 4 in the xy -plane are given different trajectories of a light ray assuming that $\frac{d\rho}{d\varphi} = 0$ at the initial point.

If the light ray cut the S.s., then it will tend to the center $r = 0$, because the equation (3.2) has no solution. The equations of motion of a light ray are simpler, because $\beta = \infty$. The equation (3.3) for light rays becomes

$$\left(\frac{d\rho}{d\varphi}\right)^2 = \left[\frac{\rho_0^2}{\left(1 + \frac{GM\rho_0}{c^2}\right)^4} - \frac{\rho^2}{\left(1 + \frac{GM\rho}{c^2}\right)^4} \right] \left(1 + \frac{GM\rho}{c^2}\right)^4, \quad (3.7)$$




Fig.4

while the equation

$$r^2 \frac{d\varphi}{dt} = C_2 \exp\left(-\int \frac{GM}{\mu c^2} \left(2 + \frac{1}{\beta(\beta+1)}\right) d\rho\right)$$

becomes

$$r^2 \frac{d\varphi}{dt} = C_2 \left(1 + \frac{GM\rho}{c^2}\right)^{-2}.$$

Assuming that $\frac{d\rho}{d\varphi} = 0$ if $r = r_0$, we find

$$C_2 = r_0^2 \frac{d\varphi}{dt} \left(1 + \frac{GM\rho_0}{c^2}\right)^2 = r_0 c \left(1 + \frac{GM\rho_0}{c^2}\right)^2,$$

and hence

$$r^2 \frac{d\varphi}{dt} = r_0 c \left(\frac{1 + \frac{GM}{r_0 c^2}}{1 + \frac{GM}{rc^2}}\right)^2. \quad (3.8)$$

The total angle of deflection of a light ray outside the S.s. is equal to

$$\Delta\alpha = 2 \int_0^{\rho_0} \frac{d\rho}{\sqrt{\frac{\rho_0^2}{(1+GM\rho_0/c^2)^4} (1 + GM\rho/c^2)^4 - \rho^2}} - \pi. \quad (3.9)$$

(ii) Assume that $C_1 < 1$. Physically it means that the particle can not leave the gravitational field because of small energy. In this case $f(\rho)$ is defined only for $\rho > \rho_*$, where ρ_* is the solution of the equation

$$1 - \ln\left(1 + \frac{GM\rho}{c^2}\right) = C_1, \quad \rho_* = \frac{c^2}{GM} \left(e^{1-C_1} - 1\right).$$

Note that $\lim_{\rho \rightarrow \rho_*+} f(\rho) = \infty$ and $\lim_{\rho \rightarrow \infty} f(\rho) = 0$. Similarly to the case (i), the condition $\frac{df}{d\rho} = 0$ is equivalent to (3.5) and further it reduces to (3.6), but now $C_1 < 1$. Now three subcases are possible.




Fig.5

(ii_a) Assume that $0,9313 < C_1 < 1$. Then the equation (3.6) has two solutions (fig. 1), ρ_c and ρ_s . The smaller solution ρ_c has just the same meaning as in the Newton's theory: Each particle on a distance $r_c = \frac{1}{\rho_c}$ will move on a circle with velocity $\sqrt{GM\rho_c}$. The larger solution ρ_s has the same property but it is a relativistic effect, because $\rho_s > 0,2637\frac{c^2}{GM}$, ($r_s < 3,7921\frac{GM}{c^2}$), and $r_s = \frac{1}{\rho_s}$ will be called again Schwarzschild radius because it separates the curves just like on fig. 4. The graphic of the function $f(\rho)$ is presented on fig. 5.

Analogously as in the previous cases we have the following conclusions.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} > f(\rho_s)$, then the equation (3.2) has no solutions, such that on the trajectory will never be $\frac{d\rho}{d\varphi} = 0$. Thus, the particle will tend to the center $r = 0$. This easily can be achieved if we throw a particle with small velocity which is almost radial motion, and then the particle will come to the center of the Earth, assuming that mass of the Earth is concentrated at a point. According to the Newton's theory the particle will move on an ellipse with large eccentricity.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} = f(\rho_s)$, then two cases are possible. If $\rho = \rho_s$, then the particle will move on a circle on the S.s., while if $\rho < \rho_s$, then the particle will tend to the center $r = 0$.

- If $f(\rho_c) < \frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} < f(\rho_s)$, then two cases are possible. If $\rho < \rho_s$, then the equation (3.3) has two solutions smaller than ρ_s and then the trajectory will be like an ellipse, analogously to the planetary orbits. The distance r to the center of the gravitation satisfies $r_1 \leq r \leq r_2$, where $\rho_1 = \frac{1}{r_1}$ and $\rho_2 = \frac{1}{r_2}$ can be found as solutions of (3.3). On the other hand, if $\rho > \rho_s$, then the particle will remain in the S.s. and will tend to the center $r = 0$.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} = f(\rho_c)$, then two cases are possible. If $\rho < \rho_s$, then the particle will move on a circle, while if $\rho > \rho_s$, the particle will remain in the S.s. and will tend to the center $r = 0$.

- If $\frac{(\frac{d\rho}{d\varphi})^2 + \rho^2}{v^2 e^I} < f(\rho_c)$, then the particle must remain in the S.s. and will tend to the center $r = 0$.

(ii_b) Assume that $C_1 = 0,9313$. Then the equation (3.6) has a double solution $\rho_s = \rho_c = 0,2637\frac{c^2}{GM}$. In this case ρ_s and ρ_c coincide, the graphic of $f(\rho)$ is given on fig. 6. The discussion is analogous as previous.

(ii_c) Assume that $C_1 < 0,9313$. Then the equation (3.2) has no solution and the particle tends to the center $r = 0$. The graphic of $f(\rho)$ is given on fig. 7.

At the end of this section we give some global conclusions in condensed form. First note that the Schwarzschild radius is defined for each constant $C_1 \geq 0,9313$, r_s belongs to the following segment

$$\frac{GM}{c^2} \leq r_s \leq \frac{1}{0,2637} \frac{GM}{c^2} = 3,7921 \frac{GM}{c^2}$$

and r_s for the light rays is equal to $\frac{GM}{c^2}$.

- If at the initial moment the particle is inside the S.s. and the equation (3.2) has a solution $\rho > \rho_s$, then the particle has never been outside the S.s., it will remain in the S.s. and will tend to the center $r = 0$.




Fig.6




Fig.7

- Suppose that at the initial moment the particle is outside the S.s. If there exists an unique solution ρ of (3.2), such that $\rho < \rho_s$, then the trajectory will look like a parabola or hyperbola and will never pass inside the S.s. If there exist two solutions ρ_1, ρ_2 , ($\rho_1, \rho_2 > \rho_s$), then the motion will look like an ellipse.

- If the equation (3.2) has no solution, then it will never be $\frac{d\rho}{d\varphi} = 0$ on the trajectory. The particle will tend to the center $r = 0$. In this case the particle can cut the S.s.

We shall explain why the other cases are not possible. If the equation (3.2) has at least one solution, the trajectory will be axially symmetric with respect to the line which connects the center $r = 0$ and the point where $\frac{d\rho}{d\varphi} = 0$. Moreover, the trajectories are "time reversible", which means that if a particle moves in one direction on the trajectory, then it can move also in the opposite direction of that trajectory.

We shall prove the following statements.

Proposition 1. *If a particle cut the S.s., then inside the S.s. can never be $\frac{d\rho}{d\varphi} = 0$, and the particle tends to the center of gravitation.*

Proof. Assume that ρ , v and $\frac{d\rho}{d\varphi}$ are given. Then ρ and v determine the value of C_1 . Now the equation (3.2) can not have a solution, because if such a solution ρ' exists, then $\rho' < \rho$, because $f(\rho)$ is a decreasing function for $\rho > \rho_s$. This finishes the proof.

Corollary 1. *The trajectories of "elliptic type" can not cut the S.s.*

Proposition 2. *Inside the S.s. there does not exist a trajectory of "elliptic type".*

Proof. Assume that there is a trajectory of elliptic type which is inside the S.s. and let the extreme distances of the particle to the center of gravitation be $r_1 = \frac{1}{\rho_1}$ and $r_2 = \frac{1}{\rho_2}$. Then there will exist a radius $r \in (r_1, r_2)$, such that for the same value of C_1 , a new particle can move on a circle of radius r . This is a contradiction with the fact that each S.s. is not contained inside another S.s. because the radius of the S.s. is minimal such that a particle can move on a circle for the same value of C_1 .

4 Some relationships between the geodesics in ILLTG and GR

In this section we shall find some connections between the planetary orbits according to ILLTG using orthonormal coordinates and GR using the Schwarzschild solution. Namely, we shall show that there are simple transformations which the geodesics from ILLTG map into the geodesics from GR and conversely.

Let us denote by r , $\rho = \frac{1}{r}$, φ , t the polar coordinates in ILLTG, and by r_* , $\rho_* = \frac{1}{r_*}$, φ_* , t_* we shall denote the polar coordinates in GR. Note that in both tests in the solar system (deflection of the light ray near the Sun and the perihelion shift) were measured angles and these angles were the same for ILLTG and GR. This associates us to assume that $\varphi_* = \varphi$, i.e. we can choose the angle φ as a common parameter, and then we can find the relationships between ρ and ρ_* , as well as between t and t_* . These relationships will be found by neglecting the expressions of order c^{-4} . We convenient to write φ instead of φ_* . Although we shall consider ellipses as planetary trajectories, the obtained results are also true if the trajectory of a planet is a hyperbola or a parabola.

Let us start with the equations

$$d\varphi = -\frac{(1 + 3\frac{GM}{ac^2(1-\epsilon^2)})d\rho}{\sqrt{(\rho_1 - \rho)(\rho - \rho_2)}}, \quad (4.1)$$

$$d\varphi = -\frac{d\rho_*}{\sqrt{(\rho_1 - \rho_*)(\rho_* - \rho_2)}} \left[1 + \frac{GM}{c^2}\rho_* + \frac{2GM}{ac^2(1-\epsilon^2)} \right] \quad (4.2)$$

from the ILLTG and GR respectively. The solution of (4.1) we found to be

$$\rho = \frac{\rho_1 + \rho_2}{2} + \frac{\rho_1 - \rho_2}{2} \cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1-\epsilon^2)} \right) \right] \quad (4.3)$$

and the solution of (4.2) is given by

$$\begin{aligned} \varphi &= -\left(1 + \frac{2GM}{ac^2(1-\epsilon^2)} \right) \left(\arcsin \frac{2\rho_* - \rho_1 - \rho_2}{\rho_1 - \rho_2} - \frac{\pi}{2} \right) + \\ &+ \frac{GM}{c^2} \left[-\sqrt{(\rho_1 - \rho_*)(\rho_* - \rho_2)} + \frac{\rho_1 + \rho_2}{2} \left(-\arcsin \frac{2\rho_* - \rho_1 - \rho_2}{\rho_1 - \rho_2} + \frac{\pi}{2} \right) \right] = \\ &= -\left(1 + \frac{3GM}{ac^2(1-\epsilon^2)} \right) \left(\arcsin \frac{2\rho_* - \rho_1 - \rho_2}{\rho_1 - \rho_2} - \frac{\pi}{2} \right) - \frac{GM}{c^2} \sqrt{(\rho_1 - \rho_*)(\rho_* - \rho_2)} \end{aligned}$$

and hence

$$\cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1-\epsilon^2)} \right) \right] = \frac{2\rho_* - \rho_1 - \rho_2}{\rho_1 - \rho_2} + \frac{GM}{c^2} \sqrt{(\rho_1 - \rho_*)(\rho_* - \rho_2)} \sqrt{1 - \left(\frac{2\rho_* - \rho_1 - \rho_2}{\rho_1 - \rho_2} \right)^2},$$

i.e.

$$\cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1-\epsilon^2)} \right) \right] = \frac{2\rho_* - \rho_1 - \rho_2}{\rho_1 - \rho_2} + \frac{GM}{c^2} \frac{2(\rho_1 - \rho_*)(\rho_* - \rho_2)}{\rho_1 - \rho_2}. \quad (4.4)$$

By eliminating the angle φ from (4.3) and (4.4), we get

$$\rho = \rho_* + \frac{GM}{c^2} (\rho_1 - \rho_*)(\rho_* - \rho_2) \quad (4.5)$$

and also the converse relation

$$\rho_* = \rho - \frac{GM}{c^2} (\rho_1 - \rho)(\rho - \rho_2). \quad (4.5')$$

Now we shall find the analogous equation of (4.1) written in ordinary coordinates. We shall use the letter R for radial distance in ordinary coordinates. First let us write the solution (4.3) in the following form

$$r = \left[\frac{\rho_1 + \rho_2}{2} + \frac{\rho_1 - \rho_2}{2} \cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1 - \epsilon^2)} \right) \right] \right]^{-1}$$

and after differentiating, using that $dr = \left(1 - \frac{GM}{rc^2} \right) dR \approx \left(1 - \frac{GM}{Rc^2} \right) dR$, we obtain

$$\left(1 - \frac{GM}{Rc^2} \right) dR = d \left\{ \left[\frac{\rho_1 + \rho_2}{2} + \frac{\rho_1 - \rho_2}{2} \cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1 - \epsilon^2)} \right) \right] \right]^{-1} \right\}.$$

After integration of this equation we get

$$R - \frac{GM}{c^2} \ln \frac{R}{R_1} = \left[\frac{\rho_1 + \rho_2}{2} + \frac{\rho_1 - \rho_2}{2} \cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1 - \epsilon^2)} \right) \right] \right]^{-1},$$

where the constant of integration is chosen such that $R = R_1 = (\rho_1)^{-1}$ for $\varphi = 0$. Further, for elliptic trajectories with small eccentricity we can put $\ln \frac{R}{R_1} \approx \frac{R}{R_1} - 1$, and hence

$$R = -\frac{GM}{c^2} + \left(1 + \frac{GM\rho_1}{c^2} \right) \left[\frac{\rho_1 + \rho_2}{2} + \frac{\rho_1 - \rho_2}{2} \cos \left[\varphi \left(1 - \frac{3GM}{ac^2(1 - \epsilon^2)} \right) \right] \right]^{-1}. \quad (4.6)$$

This function R for $\varphi = 0$ and $\varphi = \frac{\pi}{2} \left(1 + \frac{3GM}{ac^2(1 - \epsilon^2)} \right)$ takes values $(\rho_1)^{-1}$ and $(\rho_2)^{-1} \left(1 + \frac{GM\rho_1}{c^2} \right) - \frac{GM}{c^2} \approx (\rho_2)^{-1} \left(1 + \frac{GM(\rho_2 - \rho_1)}{c^2} \right)^{-1}$ respectively. Thus, in (4.6) ρ_2 should be replaced by $\rho_2 \left(1 - \frac{GM(\rho_2 - \rho_1)}{c^2} \right)$. Hence we obtain

$$R = -\frac{GM}{c^2} + \left(1 + \frac{GM\rho_1}{c^2} \right) \left\{ \frac{1}{2} \left[\rho_1 + \rho_2 - \rho_2 \frac{GM(\rho_2 - \rho_1)}{c^2} \right] + \frac{1}{2} \left[\rho_1 - \rho_2 + \rho_2 \frac{GM(\rho_2 - \rho_1)}{c^2} \right] \cos \varphi \left(1 - \frac{3GM}{ac^2(1 - \epsilon^2)} \right) \right\}^{-1}.$$

After some calculations, one can verify that the function R^{-1} satisfies the equality

$$R^{-1} = \rho - \frac{GM}{c^2} (\rho_1 - \rho)(\rho - \rho_2), \quad (4.7)$$

i.e.

$$\rho_* = R^{-1}. \quad (4.8)$$

The differential equation for R^{-1} , analogous to (4.1) is given by

$$d\varphi = -\frac{d\frac{1}{R}}{\sqrt{(\rho_1 - R^{-1})(R^{-1} - \rho_2)}} \left[1 + \frac{GM}{c^2} \frac{1}{R} + \frac{2GM}{ac^2(1 - \epsilon^2)} \right]. \quad (4.9)$$

Note that if we put $r = r_* \left(1 - \frac{GM}{r_* c^2} \right) = r_* - \frac{GM}{c^2}$, then the equation (4.1) becomes

$$d\varphi = -\frac{d\rho_*}{\sqrt{(\rho_{1*} - \rho_*)(\rho_* - \rho_{2*})}} \left[1 + \frac{GM}{c^2} \rho_* + \frac{2GM}{ac^2(1 - \epsilon^2)} \right], \quad (4.9')$$

where $\rho_{1*} = (r_1 + \frac{GM}{c^2})^{-1}$ and $\rho_{2*} = (r_2 + \frac{GM}{c^2})^{-1}$.

The equation (4.8) shows that **for $v \ll c$, the trajectory of motion in the (r, φ) plane is the same with the trajectory according ILLTG using ordinary coordinates.**

Further we shall find the relationships between t , t_* and T . According to GR it is

$$r_*^2 \frac{d\varphi}{dt_*} = C_1 \left(1 - \frac{2GM}{c^2} \rho_* \right)$$

and according to ILLTG it is

$$r^2 \frac{d\varphi}{dt} = C_2 \left(1 - \frac{2GM}{c^2} \rho - \frac{1}{\beta(\beta+1)} \frac{GM\rho}{c^2} \right).$$

The quotient of these two equations yields to

$$\left(\frac{r_*}{r} \right)^2 \frac{dt}{dt_*} = C_3 \left(1 + \frac{1}{\beta(\beta+1)} \frac{GM\rho}{c^2} \right).$$

Since $\frac{r_*}{r} = 1 + \frac{GM}{\rho c^2}(\rho_1 - \rho)(\rho - \rho_2)$ according to (4.5'), and $\frac{1}{\beta(\beta+1)} \approx \frac{1}{2}$ we obtain

$$\frac{dt}{dt_*} = C_3 \left(1 + \frac{1}{2} \frac{GM\rho}{c^2} - 2 \frac{GM}{\rho c^2} (\rho_1 - \rho)(\rho - \rho_2) \right).$$

Moreover, $C_3 = 1$, because in case of parabola or hyperbola, $\frac{dt}{dt_*} \rightarrow 1$ if $\rho \rightarrow 0$. Hence we get

$$\frac{dt}{dt_*} = 1 + \frac{1}{2} \frac{GM\rho}{c^2} - 2 \frac{GM}{\rho c^2} (\rho_1 - \rho)(\rho - \rho_2).$$

Assuming that the trajectory is elliptic with eccentricity close to 0, then

$$\frac{dt}{dt_*} = 1 + \frac{1}{2} \frac{GM\rho}{c^2}.$$

Moreover, we can find the connection between t_* and the ordinary time coordinate T from ILLTG using that $dt = \left(1 + \frac{GM\rho}{c^2} \right) dT$, i.e.

$$\frac{dT}{dt_*} = 1 - \frac{1}{2} \frac{GM\rho}{c^2}. \quad (4.10)$$

The formula (4.10) does not show how much the time according to GR is "faster" (or "slower") than the time according to ILLTG, but only a factor for transition from the geodesic of GR into geodesic of ILLTG in ordinary coordinates.

Further we shall compare also the null geodesics, i.e. the trajectories of the light rays. According to (2.14) in ILLTG it is

$$\rho = \frac{1}{R} \cos \varphi + \frac{GM}{R^2 c^2} (2 - 2 \cos \varphi), \quad (4.11)$$

while according to GR it is [5]

$$\rho_* = \frac{1}{R} \cos \varphi + \frac{GM}{R^2 c^2} (2 - \cos^2 \varphi). \quad (4.12)$$

Hence we obtain

$$\rho_* = \rho + \frac{GM}{R^2 c^2} \cos \varphi (2 - \cos \varphi), \quad (4.13)$$

$$\frac{\rho_*}{\rho} = 1 + \frac{GM}{\rho R^2 c^2} \cos \varphi (2 - \cos \varphi),$$

i.e.

$$\frac{r_*}{r} = 1 - \frac{GM r}{R^2 c^2} \cos \varphi (2 - \cos \varphi). \quad (4.14)$$

Further, using (4.14) as well as the following two equalities

$$r^2 \frac{d\varphi}{dt} = C \left(1 - 2 \frac{GM}{r c^2} \right),$$

$$r_*^2 \frac{d\varphi}{dt_*} = C \left(1 - 2 \frac{GM}{r_* c^2} \right),$$

which are almost the same in both theories, we get

$$\begin{aligned} \left(\frac{r_*}{r} \right)^2 \frac{dt}{dt_*} &= 1, \\ \frac{dt}{dt_*} &= 1 + 2 \frac{GM r}{R^2 c^2} \cos \varphi (2 - \cos \varphi). \end{aligned} \quad (4.15)$$

The angle of deflection from a point A to a point B when the ray passes near the Sun according to (2.16) in ILLTG is equal to

$$\Delta\alpha_1 = \frac{GM}{Rc^2} \left(2\sqrt{1 - \frac{R^2}{r_1^2}} + 2\sqrt{1 - \frac{R^2}{r_2^2}} \right), \quad (4.16)$$

and also using PPN coordinates GR yields to the same formula (4.16). This value (4.16) is confirmed with a very precise measurements in the recent test with Cassini spacecraft [4]. According to ILLTG, the angle of deflection is always twice larger than the classical Newton's result, and can be explained with the coefficient $2 - \beta^{-2} = 1 + v^2/c^2 = 2$ in (2.2) when $v = c$. Note that this deflection of the light is not a "gravitational effect", because it will appear if we have only an acceleration without gravitation, and ILLTG differs slightly in these two cases when there is a gravitation and there is only simulation of gravitation using inertial forces. About this will be said more at the end of this section.

We shall compare now the radial motions of a particle with large velocity v in a gravitational field. The Newton's relation $\frac{1}{2}v^2 - GM\rho = \text{const.}$ according to ILLTG should be replaced by (2.2c), i.e.

$$\frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} - \ln \left(1 + \frac{GM\rho}{c^2} \right) = \text{const.} \quad (4.17)$$

According to GR, for a radial motion in a gravitational field we get

$$\frac{\sqrt{1 - \frac{2GM\rho}{c^2}}}{\sqrt{1 - \frac{v^2}{c_*^2}}} = \text{const.} \quad (4.18)$$

where $c_* = c(1 - \frac{2GM\rho}{c^2})$ is the velocity of the light measured in the gravitational field. The physical interpretation for both formulas (4.17) and (4.18) obtains if we multiply them by $m_0 c^2$. From (4.17) we obtain

$$E_k + E_p = \text{const.} \quad (4.19)$$

because in section 6 will be proved that the potential energy for a particle with small mass is equal to $E_p = -m_0 c^2 \ln \left(1 + \frac{GM\rho}{c^2} \right)$. The equation (4.18) can be written as

$$E_{\text{total}} \sqrt{1 - \frac{2GM\rho}{c^2}} = \text{const.} \quad (4.20)$$

If $\frac{GM\rho}{c^2}$ is very small, i.e. we consider a weak gravitational field, (4.18) can be written in the form

$$\frac{1}{\sqrt{1 - \frac{v^2}{c_*^2}}} = C \frac{GM\rho}{c^2} + C, \quad C = \text{const.}$$

and hence

$$\frac{d}{dt} \frac{1}{\sqrt{1 - \frac{v^2}{c_*^2}}} = C \frac{GM}{c^2} \frac{d}{dt} \frac{1}{r},$$

i.e.

$$\frac{d}{dt} \frac{1}{\sqrt{1 - \frac{v^2}{c_*^2}}} = C \frac{a}{c^2} v, \quad (4.21)$$

where $a = -\frac{GM}{r^2}$ is the acceleration and $v = dr/dt$. The analogous equation from (4.17) in weak gravitational field is the following

$$\frac{d}{dt} \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} = \frac{a}{c^2} v. \quad (4.22)$$

In a short time interval the acceleration a is almost constant and the equations (4.21) and (4.22) should be close to the equation of motion of a particle in STR under a constant acceleration a . The STR motion

$$\frac{d}{\sqrt{1 - \frac{v^2}{c^2}} dt} \frac{m_0 c^2}{\sqrt{1 - \frac{v^2}{c^2}}} = a \frac{m_0 v}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (4.23)$$

is just the equation (4.22). Indeed, in (4.23) a is a constant, because the acceleration a is unchanged under the Lorentz transformations of motion in the x direction, because the tensor with components

$$\begin{bmatrix} 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ -a & 0 & 0 & 0 \end{bmatrix}$$

is unchanged under such Lorentz transformations, analogously as the component E_x of the electric field is constant under such Lorentz transformations.

Both of equations (4.21) and (4.22) in the (xt) -plane are hyperbolas, but with different parameters. The parameter C has no physical meaning according to STR for a motion under a constant acceleration, although this constant in (4.18) has a good physical meaning. Thus, the equation of radial motion is not close to the STR. Moreover, from (4.18) one can conclude that $\frac{dv}{dt} \approx a(1 - 3\frac{v^2}{c^2})$, such that the acceleration changes its sign if $v > c/\sqrt{3}$.

We note that ILLTG is very close to the STR. Namely, if we consider a noninertial system without gravitation but there is only a 3-vector of acceleration (a_x, a_y, a_z) and a 3-vector of angular velocity (w_x, w_y, w_z) , then in the equations of motion (2.3) we should only put $\mu = 1$. This is obtained in the frame of STR, under the assumption that the Jacobi matrices with respect to the inertial systems are orthogonal and hence the coordinates in such a system are unholonomic [9]. The influence of the coefficient μ in (2.2) and (2.3) is very small. For example, if we have a noninertial system without gravitation with the same acceleration $(a_x, a_y, a_z) = \left(-\frac{GM}{r^3}x, -\frac{GM}{r^3}y, -\frac{GM}{r^3}z\right)$, then the angle of deflection of a light ray near the Sun will be again $\Delta\alpha = \frac{4GM}{Rc^2}$, while the angle between two perihelions will be $\Delta\alpha = \frac{7GM\pi}{ac^2(1-\epsilon^2)}$, instead of $\Delta\alpha = \frac{6GM\pi}{ac^2(1-\epsilon^2)}$. The equation (2.4) will not be changed, and the equation (4.17) will be replaced by

$$\frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} - \frac{GM\rho}{c^2} = const.$$

5 Two proposal tests

After the discussion in section 4 we propose the following tests for which we have different results from ILLTG and GR.

Test 1. Let A_1, A_2 and A_3 are three not close points of a planet orbit or a spacecraft trajectory, and let $\angle A_0OA_1 = \varphi_0$, $\angle A_0OA_2 = \varphi_1$, $\angle A_0OA_3 = \varphi_2$, where A_0 is the perihelion of the planet orbit and O is the center of the Sun. We denote by T_1 and T_2 the times for which the planet comes from

A_1 to A_2 and from A_2 to A_3 respectively. We will calculate $T_1 : T_2$ according to ILLTG and GR, and later can be compared with the experimental results. We shall find $T_1 : T_2$ where T_1 and T_2 are the proper times for the Earth, where the trajectory of the Earth is observed. Let us denote the proper time for the Earth by τ .

First let us calculate $T_1 : T_2$ according to ILLTG. From (2.4) we obtain

$$r^2 \frac{d\varphi}{d\tau} = r^2 \frac{d\varphi}{dt} \frac{dt}{dT} \frac{dT}{d\tau} = C_2 \left(1 - \frac{5}{2} \frac{GM}{rc^2}\right) \left(1 + \frac{GM}{rc^2}\right) \left(1 + \frac{GM}{rc^2}\right) = C_2 \left(1 - \frac{1}{2} \frac{GM}{rc^2}\right).$$

Thus

$$d\tau = \frac{d\varphi}{\rho^2 C_2 (1 - \frac{1}{2} \frac{GM\rho}{c^2})} = \frac{1}{C_2} \rho^{-2} \left(1 + \frac{1}{2} \frac{GM\rho}{c^2}\right) d\varphi,$$

and hence

$$T_1 : T_2 = \int_{\varphi_0}^{\varphi_1} \left(\rho^{-2} + \frac{1}{2} \frac{GM\rho^{-1}}{c^2}\right) d\varphi : \int_{\varphi_1}^{\varphi_2} \left(\rho^{-2} + \frac{1}{2} \frac{GM\rho^{-1}}{c^2}\right) d\varphi,$$

where ρ is given by (4.3). Using that $\frac{GM}{c^2}$ is close to 0, and replacing $\varphi \left(1 - \frac{3GM}{ac^2(1-\epsilon^2)}\right)$ with a new variable, one obtains

$$T_1 : T_2 = \frac{A}{C} - \frac{3GM}{ac^2(1-\epsilon^2)} \frac{C(\rho^{-2}(\varphi_1)\varphi_1 - \rho^{-2}(\varphi_0)\varphi_0) - A(\rho^{-2}(\varphi_2)\varphi_2 - \rho^{-2}(\varphi_1)\varphi_1)}{C^2} + \frac{1}{2} \frac{GM}{c^2} \frac{BC - AD}{C^2}, \quad (5.1)$$

where

$$A = \int_{\varphi_0}^{\varphi_1} \rho^{-2} d\varphi, \quad B = \int_{\varphi_0}^{\varphi_1} \rho^{-1} d\varphi, \quad C = \int_{\varphi_1}^{\varphi_2} \rho^{-2} d\varphi, \quad D = \int_{\varphi_1}^{\varphi_2} \rho^{-1} d\varphi.$$

After solving these integrals, where $\rho = \frac{1}{2}[(\rho_1 + \rho_2) + (\rho_1 - \rho_2) \cos \varphi]$, we obtain

$$A = -\frac{\rho_1 + \rho_2}{(\rho_1 \rho_2)^{3/2}} \arctan \frac{\sqrt{\rho_1 \rho_2} (\cot \frac{\varphi_1}{2} - \cot \frac{\varphi_0}{2})}{\rho_2 + \rho_1 \cot \frac{\varphi_1}{2} \cot \frac{\varphi_0}{2}} + \frac{\rho_2 - \rho_1}{\rho_1 \rho_2} \left(\frac{\cot \frac{\varphi_1}{2}}{\rho_2 + \rho_1 \cot^2 \frac{\varphi_1}{2}} - \frac{\cot \frac{\varphi_0}{2}}{\rho_2 + \rho_1 \cot^2 \frac{\varphi_0}{2}} \right), \quad (5.2)$$

$$C = -\frac{\rho_1 + \rho_2}{(\rho_1 \rho_2)^{3/2}} \arctan \frac{\sqrt{\rho_1 \rho_2} (\cot \frac{\varphi_2}{2} - \cot \frac{\varphi_1}{2})}{\rho_2 + \rho_1 \cot \frac{\varphi_2}{2} \cot \frac{\varphi_1}{2}} + \frac{\rho_2 - \rho_1}{\rho_1 \rho_2} \left(\frac{\cot \frac{\varphi_2}{2}}{\rho_2 + \rho_1 \cot^2 \frac{\varphi_2}{2}} - \frac{\cot \frac{\varphi_1}{2}}{\rho_2 + \rho_1 \cot^2 \frac{\varphi_1}{2}} \right), \quad (5.3)$$

$$B = \frac{2}{\sqrt{\rho_1 \rho_2}} \arctan \frac{\sqrt{\rho_1 \rho_2} (\tan \frac{\varphi_1}{2} - \tan \frac{\varphi_0}{2})}{\rho_1 + \rho_2 \tan \frac{\varphi_1}{2} \tan \frac{\varphi_0}{2}}, \quad (5.4)$$

$$D = \frac{2}{\sqrt{\rho_1 \rho_2}} \arctan \frac{\sqrt{\rho_1 \rho_2} (\tan \frac{\varphi_2}{2} - \tan \frac{\varphi_1}{2})}{\rho_1 + \rho_2 \tan \frac{\varphi_2}{2} \tan \frac{\varphi_1}{2}}. \quad (5.5)$$

Now let us calculate $T_1 : T_2$ according to GR. Using that $r_*^2 \frac{d\varphi}{d\tau_*} = C_1$, and (4.5') we obtain

$$d\tau_* = \frac{d\varphi}{\rho_*^2 C_1} = \frac{d\varphi}{C_1 \rho^2 \left[1 - \frac{2GM\rho}{c^2} \left(\frac{\rho_1}{\rho} - 1\right) \left(1 - \frac{\rho_2}{\rho}\right)\right]},$$

$$\tau_* = \frac{1}{C_1} \int \rho^{-2} \left[1 + \frac{2GM\rho}{c^2} \left(\frac{\rho_1}{\rho} - 1\right) \left(1 - \frac{\rho_2}{\rho}\right)\right] d\varphi,$$

where ρ is given by (4.3). If the eccentricity of the ellipse is small, then the expression $\frac{2GM\rho}{c^2} \left(\frac{\rho_1}{\rho} - 1\right) \left(1 - \frac{\rho_2}{\rho}\right)$ can be neglected. Then, analogously to (5.1) we obtain

$$T_1 : T_2 = \frac{A}{C} - \frac{3GM}{ac^2(1-\epsilon^2)} \frac{C(\rho^{-2}(\varphi_1)\varphi_1 - \rho^{-2}(\varphi_0)\varphi_0) - A(\rho^{-2}(\varphi_2)\varphi_2 - \rho^{-2}(\varphi_1)\varphi_1)}{C^2} \quad (5.6)$$

where A , C , B and D are given by (5.2-5).

It is convenient to assume that $\varphi_0 = 0$, $\varphi_1 = \pi/2$ and $\varphi_2 = \pi$. Then

$$A = \frac{\rho_1 + \rho_2}{(\rho_1 \rho_2)^{3/2}} \arctan \sqrt{\frac{\rho_2}{\rho_1}} + \frac{\rho_2 - \rho_1}{(\rho_1 + \rho_2) \rho_1 \rho_2}, \quad C = \frac{\rho_1 + \rho_2}{(\rho_1 \rho_2)^{3/2}} \arctan \sqrt{\frac{\rho_1}{\rho_2}} - \frac{\rho_2 - \rho_1}{(\rho_1 + \rho_2) \rho_1 \rho_2}, \quad (5.7)$$

$$B = \frac{2}{\sqrt{\rho_1 \rho_2}} \arctan \sqrt{\frac{\rho_2}{\rho_1}}, \quad D = \frac{2}{\sqrt{\rho_1 \rho_2}} \arctan \sqrt{\frac{\rho_1}{\rho_2}}. \quad (5.8)$$

Remember that both formulas (5.1) and (5.6) are obtained for the observers on the planet whose trajectory is observed. Indeed, it means that these formulas are valid only for the trajectory of the Earth, because we observe from the Earth. Further let us modify these two formulas assuming that the observer is far from gravitation in the solar system. Then, according to ILLTG we obtain

$$dt = \frac{d\varphi(1 + 2\frac{GM\rho}{c^2})}{\rho^2 C_2(1 - \frac{1}{2}\frac{GM\rho}{c^2})} = \frac{1}{C_2} \rho^{-2} \left(1 + \frac{5}{2} \frac{GM\rho}{c^2}\right) d\varphi,$$

$$T_1 : T_2 = \int_{\varphi_0}^{\varphi_1} \left(\rho^{-2} + \frac{5}{2} \frac{GM\rho^{-1}}{c^2}\right) d\varphi : \int_{\varphi_1}^{\varphi_2} \left(\rho^{-2} + \frac{5}{2} \frac{GM\rho^{-1}}{c^2}\right) d\varphi,$$

and hence

$$T_1 : T_2 = \frac{A}{C} - \frac{3GM}{ac^2(1 - \epsilon^2)} \frac{C(\rho^{-2}(\varphi_1)\varphi_1 - \rho^{-2}(\varphi_0)\varphi_0) - A(\rho^{-2}(\varphi_2)\varphi_2 - \rho^{-2}(\varphi_1)\varphi_1)}{C^2} +$$

$$+ \frac{5}{2} \frac{GM}{c^2} \frac{BC - AD}{C^2}, \quad (5.1')$$

where A , B , C and D were previously determined. Analogously, according to GR instead of (5.6) we have the formula

$$T_1 : T_2 = \frac{A}{C} - \frac{3GM}{ac^2(1 - \epsilon^2)} \frac{C(\rho^{-2}(\varphi_1)\varphi_1 - \rho^{-2}(\varphi_0)\varphi_0) - A(\rho^{-2}(\varphi_2)\varphi_2 - \rho^{-2}(\varphi_1)\varphi_1)}{C^2} +$$

$$+ 2 \frac{GM}{c^2} \frac{BC - AD}{C^2}, \quad (5.6')$$

where A , C , B and D were previously determined.

Finally, if we measure from the Earth the trajectory of another planet, then we should modify the formulas (5.1') and (5.6') because the trajectory of the Earth is not a circle, the "speed" of the time is proportional to $(1 - \frac{GM}{rc^2})^2$, where r is the distance from the Earth to the center of the Sun. Thus, let us denote by ν_1 and ν_2 the average values of the "speed" of the time on the Earth during the first and the second time interval. Thus, on the Earth we shall measure the quotient

$$\frac{\nu_1 T_1}{\nu_2 T_2} = \frac{\nu_1}{\nu_2} \frac{T_1}{T_2} \quad (5.9)$$

where $T_1 : T_2$ is given by (5.1') and (5.6') respectively according to ILLTG and GR.

Note that in this test we should measure only the quotient of long time intervals, large angles and we need also the quotient $\rho_1 : \rho_2 = (1 + \epsilon) : (1 - \epsilon)$. The perturbations from the other planets, Moon, stars, galaxy and universe (see sec. 8 and Ref. [10,11]) are not included in the final results.

Test 2. Let us consider the second Kepler's law, i.e. consider the expression $r^2 \frac{d\varphi}{d\tau}$ on the trajectory, where τ is the proper time. We shall present the formulas assuming that we observe the trajectory of the Earth from the Earth.

According to ILLTG we have

$$r^2 \frac{d\varphi}{d\tau} = r^2 \frac{d\varphi}{dt} \frac{dt}{dT} \frac{dT}{d\tau} = C' \left(1 - \frac{5}{2} \frac{GM}{rc^2}\right) \left(1 + \frac{GM}{rc^2}\right) \left(1 + \frac{GM}{rc^2}\right) = C' \left(1 - \frac{1}{2} \frac{GM}{rc^2}\right)$$

and according to GR (omitting the star indices), it is

$$r^2 \frac{d\varphi}{d\tau} = C'',$$

where C' and C'' are constants close to $C = \sqrt{GMa(1-\epsilon^2)}$. Let P and Q be two points on the orbit and let us denote the corresponding distances to the center of the Sun by r_1 and r_2 respectively. For example, P can be the perihelion of the Earth and Q aphelion. Using the previous formulas according to ILLTG we obtain

$$r_2^2 \left(\frac{d\varphi}{d\tau} \right)_Q - r_1^2 \left(\frac{d\varphi}{d\tau} \right)_P = C \left(-\frac{1}{2} \frac{GM}{r_2 c^2} + \frac{1}{2} \frac{GM}{r_1 c^2} \right) = \frac{1}{2} \frac{GM(r_2 - r_1)}{r_1 r_2 c^2} \sqrt{GMa(1-\epsilon^2)} \quad (5.10)$$

and according to GR we obtain

$$r_2^2 \left(\frac{d\varphi}{d\tau} \right)_Q - r_1^2 \left(\frac{d\varphi}{d\tau} \right)_P = 0. \quad (5.11)$$

The perturbations from the other gravitational objects are not included in (5.10) and (5.11).

Note that in the tests 1 and 2 the departure between the GR and ILLTG is proportional to $\frac{1}{2} \frac{GM}{rc^2}$. The coefficient $\frac{1}{2}$ here is just $\frac{1}{\beta(\beta+1)}$.

6 Mass and energy in gravitational field according to ILLTG

If a particle with mass m , measured when it is far from gravitational field, has come in a gravitational field with potential $\mu = 1 + \frac{GM}{rc^2}$, then its mass measured from an observer far from gravitation, approximately is equal to $\frac{m}{1 + \frac{GM}{rc^2}}$. We convenient in this section that $M, m_1, m_2, \dots$ will mean proper mass, measured when both the observer and the mass are far from gravitation.

Let us consider two particles with masses m_1 and m_2 . Assume that the distance between their centers is R . Then the gravitational force between them is given approximately by

$$f = G \frac{m_1}{1 + \frac{Gm_2}{Rc^2}} \frac{m_2}{1 + \frac{Gm_1}{Rc^2}} \frac{1}{R^2}, \quad (6.1)$$

because each of the masses changes under the gravitational field of the other particle. This formula becomes more precise if one of the masses is very close to zero, but (6.1) is of sufficient accuracy. For example, if $m_1 \approx 0$ is mass of a small particle and $m_2 = M$ is the mass of the Sun, then $f = \frac{\frac{GmM}{R^2}}{1 + \frac{GM}{Rc^2}}$ and the acceleration for the particle is $a = \frac{\frac{GM}{R^2}}{1 + \frac{GM}{Rc^2}}$, which is in accordance with the formulas in section 2.

The energy which needs to distinguish both objects on a large distance is equal to

$$\begin{aligned} E &= \int_R^\infty f dr = \int_R^\infty G \frac{m_1}{1 + \frac{Gm_2}{rc^2}} \frac{m_2}{1 + \frac{Gm_1}{rc^2}} \frac{1}{r^2} dr, \\ E &= \frac{m_1 m_2 c^2}{m_1 - m_2} \left[\ln \left(1 + \frac{Gm_1}{Rc^2} \right) - \ln \left(1 + \frac{Gm_2}{Rc^2} \right) \right], \end{aligned} \quad (6.2)$$

which corresponds to the defect of mass

$$\Delta m = \frac{m_1 m_2}{m_1 - m_2} \left[\ln \left(1 + \frac{Gm_1}{Rc^2} \right) - \ln \left(1 + \frac{Gm_2}{Rc^2} \right) \right]. \quad (6.3)$$

If we assume in (6.2) that $m_1 = M$ is a mass of a large body for example a star, and $m_2 = m$ has negligible mass compared with M , then according to (6.3) for the potential energy of the particle we get

$$E_p = -mc^2 \ln \left(1 + \frac{GM}{Rc^2} \right). \quad (6.4)$$

Later we will return to this formula. In opposite case, if $m_1 = m_2 = m$, analogously to (6.2) the potential energy is equal to

$$E_p = -\frac{Gm^2}{R + \frac{Gm}{c^2}} \quad (6.5)$$

and the corresponding defect of mass is equal to

$$\Delta m = \frac{Gm^2}{Rc^2 + Gm}. \quad (6.6)$$

This energy can be written in the form

$$E_p = \frac{mc^2}{1 + \frac{Gm}{Rc^2}} - mc^2, \quad (6.7)$$

which is analogous to the formula for the kinetic energy according to the STR.

Further, let us consider the case when $R \rightarrow 0$. Namely, we shall consider an "ideal fusion", assuming that both particles are concentrated into points and assuming that R tends to 0. From (6.1) it just follows that the force tends to

$$f = \frac{c^4}{G}, \quad (6.8)$$

while the acceleration for each particle is of type

$$a \sim \frac{c^2}{R}. \quad (6.9)$$

Note that this acceleration does not depend on each of the masses. Probably it does not depend on the nature of these particles, for example their electricity if they are charged. In this case, when $R \rightarrow 0$, according to (6.2) for the defect of the mass we obtain

$$\Delta m = \frac{m_1 m_2}{m_1 - m_2} \ln \frac{m_1}{m_2}, \quad (6.10)$$

such that the mass after an ideal fusion is equal to

$$m = m_1 + m_2 - \frac{m_1 m_2}{m_1 - m_2} \ln \frac{m_1}{m_2}. \quad (6.11)$$

Using the differential calculus, it is easy to obtain the following inequalities:

$$m_2 \leq \frac{m_1 m_2}{m_1 - m_2} \ln \frac{m_1}{m_2} \leq \frac{m_1 + m_2}{2} \leq m_1, \quad (6.12)$$

assuming that $m_2 \leq m_1$. Specially, if $m_1 = m_2 = m$, then according to (6.6), the defect of mass is equal to $\Delta m = m$. Hence the conclusion that in "ideal fusion" we get the largest quantity of energy if $m_1 = m_2 = m$. Note also that if $m_2 \rightarrow 0$ in (6.12), then $\frac{m_1 m_2}{m_1 - m_2} \ln \frac{m_1}{m_2} \rightarrow 0$.

Analogously to the inequalities (6.12), in general case one can prove the following inequalities

$$\begin{aligned} m_2 - \frac{m_2}{1 + \frac{Gm_1}{rc^2}} &\leq \frac{m_1 m_2}{m_1 - m_2} \ln \frac{1 + \frac{Gm_1}{rc^2}}{1 + \frac{Gm_2}{rc^2}} \leq \\ &\leq \frac{1}{2} \left[\left(m_1 - \frac{m_1}{1 + \frac{Gm_2}{rc^2}} \right) + \left(m_2 - \frac{m_2}{1 + \frac{Gm_1}{rc^2}} \right) \right] \leq m_1 - \frac{m_1}{1 + \frac{Gm_2}{rc^2}}, \end{aligned} \quad (6.13)$$

if $m_2 \leq m_1$. Indeed, in special case if we put $r = 0$, we obtain the inequalities (6.12).

According to (6.4) the equation (4.17) can be written as

$$E_k + E_p = \text{const}. \quad (6.14)$$

where E_k is the kinetic energy. The equation (6.4) was obtained under the assumption that the mass of the particle is almost 0. Now, if we want to obtain the equations of motion using that the mass of the particle is not almost 0, for example motion of binary stars, we should use the potential

$$U = -\left(1 + \frac{3Gm}{rc^2}\right) \frac{c^2}{1 - \frac{m}{M}} \ln \frac{1 + \frac{GM}{rc^2}}{1 + \frac{Gm}{rc^2}}, \quad (6.15)$$

which is obtained from (6.2), where m and M are the masses of the two bodies. The "correcting multiplier" $1 + \frac{3Gm}{rc^2}$ appears if we want to use this potential to derive the equations of motion for the body with mass m . Indeed, the body with nonzero mass m influences on the gravitational potential of the other body. This potential (6.15) is sufficient to derive the approximation of the first post-Newtonian order (sec. 7). At the end of section 7 will be proved that the "correcting multiplier" must be $1 + \frac{3Gm}{rc^2}$.

If we consider the motion of a body with nonzero mass, under the gravitation of k bodies, then we can use the equations (2.19a,b,c,d), but the potential U_i instead of (2.17) is given by

$$U_i = -\left(1 + \frac{3Gm}{r_i c^2}\right) \frac{c^2}{1 - \frac{m}{m_i}} \ln \frac{1 + \frac{Gm_i}{r_i c^2}}{1 + \frac{Gm}{r_i c^2}}, \quad (6.16)$$

where m is the mass of the moving test body.

Note that (2.19) does not consider the interaction among the given k bodies, but only assumes that their positions and velocities are given. But it does not matter, because if it is given a system of n bodies with masses $m_1, m_2, \dots, m_n$, with initial positions and initial velocities, then we can find the coordinates of all bodies as functions of t by considering a system of $3n + n$ equations, i.e. 4 equations (2.19a,b,c,d) for each body, where the potential for each pair of bodies is given analogously to (6.16). Indeed, then we have the following system

$$\begin{aligned} \frac{d^2 x_j}{dt^2} = & \sum_{i=1, i \neq j}^n \left[\frac{\partial U_{ij}}{\partial x_j} \left(\beta_j^{-2} - 2 + \frac{2}{c^2} \left(\frac{dx_i}{dt} \frac{dx_j}{dt} + \frac{dy_i}{dt} \frac{dy_j}{dt} + \frac{dz_i}{dt} \frac{dz_j}{dt} \right) \right) + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial x_j} \left(M \left(\frac{dx_j}{dt} \right)^2 - 2 \frac{dx_i}{dt} \frac{dx_j}{dt} \right) \right. \\ & \left. + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial y_j} \left(M \frac{dx_j}{dt} \frac{dy_j}{dt} - 2 \frac{dx_i}{dt} \frac{dy_j}{dt} \right) + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial z_j} \left(M \frac{dx_j}{dt} \frac{dz_j}{dt} - 2 \frac{dx_i}{dt} \frac{dz_j}{dt} \right) \right] \beta_i, \quad 1 \leq j \leq n, \end{aligned} \quad (6.17a)$$

$$\begin{aligned} \frac{d^2 y_j}{dt^2} = & \sum_{i=1, i \neq j}^n \left[\frac{\partial U_{ij}}{\partial y_j} \left(\beta_j^{-2} - 2 + \frac{2}{c^2} \left(\frac{dx_i}{dt} \frac{dx_j}{dt} + \frac{dy_i}{dt} \frac{dy_j}{dt} + \frac{dz_i}{dt} \frac{dz_j}{dt} \right) \right) + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial x_j} \left(M \frac{dx_j}{dt} \frac{dy_j}{dt} - 2 \frac{dy_i}{dt} \frac{dx_j}{dt} \right) \right. \\ & \left. + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial y_j} \left(M \left(\frac{dy_j}{dt} \right)^2 - 2 \frac{dy_i}{dt} \frac{dy_j}{dt} \right) + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial z_j} \left(M \frac{dy_j}{dt} \frac{dz_j}{dt} - 2 \frac{dy_i}{dt} \frac{dz_j}{dt} \right) \right] \beta_i, \quad 1 \leq j \leq n, \end{aligned} \quad (6.17b)$$

$$\begin{aligned} \frac{d^2 z_j}{dt^2} = & \sum_{i=1, i \neq j}^n \left[\frac{\partial U_{ij}}{\partial z_j} \left(\beta_j^{-2} - 2 + \frac{2}{c^2} \left(\frac{dx_i}{dt} \frac{dx_j}{dt} + \frac{dy_i}{dt} \frac{dy_j}{dt} + \frac{dz_i}{dt} \frac{dz_j}{dt} \right) \right) + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial x_j} \left(M \frac{dx_j}{dt} \frac{dz_j}{dt} - 2 \frac{dz_i}{dt} \frac{dx_j}{dt} \right) \right. \\ & \left. + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial y_j} \left(M \frac{dy_j}{dt} \frac{dz_j}{dt} - 2 \frac{dz_i}{dt} \frac{dy_j}{dt} \right) + \frac{1}{c^2} \frac{\partial U_{ij}}{\partial z_j} \left(M \left(\frac{dz_j}{dt} \right)^2 - 2 \frac{dz_i}{dt} \frac{dz_j}{dt} \right) \right] \beta_i, \quad 1 \leq j \leq n, \end{aligned} \quad (6.17c)$$

$$\frac{d}{dt} \beta_j + \frac{1}{c^2} \sum_{i=1, i \neq j}^n \beta_i \left(\frac{\partial U_{ij}}{\partial x_j} \frac{dx_j}{dt} + \frac{\partial U_{ij}}{\partial y_j} \frac{dy_j}{dt} + \frac{\partial U_{ij}}{\partial z_j} \frac{dz_j}{dt} \right) = 0, \quad 1 \leq j \leq n, \quad (6.17d)$$

where (x_j, y_j, z_j) are the coordinates of the j -th body, $\beta_j = \left(1 - \frac{1}{c^2} \left(\left(\frac{dx_j}{dt} \right)^2 + \left(\frac{dy_j}{dt} \right)^2 + \left(\frac{dz_j}{dt} \right)^2 \right) \right)^{-1/2}$, and

$$U_{ij} = -\frac{c^2 m_i}{m_i - m_j} \ln \frac{1 + \frac{Gm_i}{r_{ij} c^2}}{1 + \frac{Gm_j}{r_{ij} c^2}} \prod_{s \neq i} \left(1 + \frac{3Gm_s}{r_{sj} c^2} \right), \quad r_{ij} = \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2}. \quad (6.18)$$

Note that $\frac{\partial U_{ij}}{\partial x_j} \frac{dx_j}{dt} + \frac{\partial U_{ij}}{\partial y_j} \frac{dy_j}{dt} + \frac{\partial U_{ij}}{\partial z_j} \frac{dz_j}{dt}$ is not a total differential like in (2.19d).

In the general equations of motion (6.17) are not included the multiple moments, the gravitational radiation and the decrease of the orbit periods of the binary systems. Indeed, these equations lead only to the first post-Newtonian order, because of the weak choice of the potentials (6.18). But, using the ordinary coordinates we have the same calculations like in GR about the gravitational radiation and the decrease of the energy and orbit period of the binary systems.

At the end of this section we shall make some considerations about the atoms of hydrogen and deuterium. First we accept the following convention. If we have a particle with mass m and electricity e , then we convert this electricity into mass $\pm \frac{e}{\sqrt{4\pi\epsilon_0 G}}$ (the sign depends on the charge positive or negative), such that the Coulon's force $f = \frac{1}{4\pi\epsilon_0} \frac{e_1 e_2}{r^2}$ can be written as a gravitational force $f = G \frac{m_1 m_2}{r^2}$. This mass will be called "charged mass". In order to distinguish the charged mass from the ordinary mass we can write $m \pm i \frac{e}{\sqrt{4\pi\epsilon_0 G}}$, where i is the imaginary unit. According to this identification, the mass of the proton is equal to

$$m_p = (1,672415 \times 10^{-24} + i \times 1,8594077 \times 10^{-6})\text{g}.$$

Let us consider the fusion of two atoms of hydrogen into atom of deuterium. According to the previous discussion, we will obtain the energy

$$E = 2 \frac{e^2 G}{Rc^2 + eG} c^2 \approx 2 \frac{e^2 G}{R},$$

where $e = 1,8594077 \times 10^{-6}\text{g}$. If we assume that $R = 2 \times 10^{-13}\text{cm}$ is the distance between the centers of the hydrogen cores just before the fusion, we get energy $E = 2,5670184 \times 10^{-27} \text{ g} \times c^2$. On the other hand, it is known from the nuclear physics that this energy is about $E = 2,569 \times 10^{-27} \text{ g} \times c^2$.

Further let us consider the question for calculation of the coefficient μ inside the atom. First we should remember that in case of gravitation it is $\mu = 1 + \frac{GM}{rc^2}$, and remember that $\frac{GM}{r} = v^2$ if the particle moves on a circle with velocity v . Hence it is better to write $\mu = 1 + \frac{v^2}{c^2}$. Now, let us consider the potential μ , when an electron is moving around the core of the atom. If its velocity is approximately $v = c/100$, then $\mu \approx 1 + 100^{-2} = 1,0001$. If we consider another particle instead of electron, for example a non charged particle, since the gravitational force is much smaller than the Coulon's force, its velocity around the core will be much smaller and hence μ will be almost 1. Hence the conclusion that *the potential μ depends on the nature of the particle*, i.e. it is not the same for all particles as it was in the previous sections.

Let us calculate the potential μ for the proton near the core of the hydrogen on a distance $2 \times 10^{-13}\text{cm}$. We can not use the formula $\mu = 1 + \frac{Ge}{Rc^2}$ where e is charged mass, but should use the following revised formula

$$\mu = 1 + \frac{Ge}{Rc^2} \lambda, \quad (6.19)$$

where the coefficient λ should be such that $\frac{Gm}{R} = v^2$. It is easy to see that the coefficient λ should be the quotient between the charged mass and the gravitational mass, i.e.

$$\lambda = \frac{1,8594077 \times 10^{-6}}{1,672415 \times 10^{-24}} \approx 1,11181 \times 10^{18}.$$

Hence, if we replace in (6.19) that $e = 1,8594077 \times 10^{-6}\text{g}$, because the ordinary mass is negligible, and $R = 2 \times 10^{-13}\text{cm}$, we get $\mu = 1,0007674584$.

If we calculate the coefficient μ for the electron close to the core, then λ will be 1836,12 times larger than the previous, because the mass of the electron is 1836,12 times smaller than the mass of the proton, while they have the same charge. In this case, for an electron on distance $2,81794 \times 10^{-13}\text{cm}$, i.e. radius of the electron, we obtain $\mu = 2$ and $\mu = 2$ achieves just for the Schwarzschild radius for the light ray (sec. 3).

Now, let us return to the potential μ for a proton near the core of the hydrogen. Since the both masses are equal, the energy obtained by fusion is equal to

$$2\left(m - \frac{m}{\mu}\right)c^2 = 2 \times 1,672415 \times 10^{-24} \text{g} \times \frac{\mu - 1}{\mu} c^2 = 2,56505 \times 10^{-27} \text{g} \times c^2.$$

7 Periastron shift of the binary systems

In this section will be calculated the periastron shift of arbitrary binary system. For example, we shall assume that it is a binary system of a pulsar and its companion. Applying the equations of motion (6.17), or simply the equations (2.18) or (2.19) with the potential (6.15) or (6.18), we shall obtain the required periastron shift.

Let m be the mass of the pulsar and M be the mass of its companion. Let us choose a coordinate system having origin at their barycenter, in the moment when they are at the periastron. Let $(x, y, 0)$ be the coordinates of the pulsar, and let $(x^*, y^*, 0)$ be the coordinates of its companion. The gravitational potential at $(x, y, 0)$ caused by the companion is given by (sec. 6)

$$U = -\left(1 + \frac{3Gm}{Rc^2}\right) \frac{c^2 M}{M - m} \ln \frac{1 + \frac{GM}{Rc^2}}{1 + \frac{Gm}{Rc^2}}, \quad (7.1)$$

where $R = \sqrt{(x - x^*)^2 + (y - y^*)^2}$. Since the following potentials

$$U' = -\left(1 + \frac{3Gm}{Rc^2}\right) \frac{c^2 M}{M + m} \ln\left(1 + \frac{G(M + m)}{Rc^2}\right) \quad (7.1')$$

and

$$U'' = -\frac{GM}{R} + \frac{1}{2} \frac{G^2 M(M - 5m)}{R^2 c^2}, \quad (7.1'')$$

do not differ from U up to c^{-2} , the same result obtains up to c^{-2} if we use alternatively the potential U' or U'' .

Let us denote the 4-vector of velocity of the pulsar by

$$(V_i) = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \left(\frac{v_x}{ic}, \frac{v_y}{ic}, \frac{v_z}{ic}, 1 \right),$$

and the 4-vector of velocity of its companion by

$$(U_i) = \frac{1}{\sqrt{1 - \frac{u^2}{c^2}}} \left(\frac{u_x}{ic}, \frac{u_y}{ic}, \frac{u_z}{ic}, 1 \right).$$

Using the potential (7.1') we can calculate the components $a_x, a_y, a_z, w_x, w_y, w_z$, which appear in (2.18). It is convenient to use the notation $r = \sqrt{x^2 + y^2}$ and $r \approx \frac{M}{M+m} R$. Let us put $\cos \alpha = \frac{x - x^*}{R}$ and $\sin \alpha = \frac{y - y^*}{R}$. Then, up to c^{-2} we have

$$\begin{aligned} a_x &= \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{\partial U}{\partial x} = \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M+m)R}{c^2}} \frac{1}{1 - 6\frac{Gm}{Rc^2}} \frac{x - x^*}{\sqrt{(x - x^*)^2 + (y - y^*)^2}} = \\ &= \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M-5m)R}{c^2}} \cos \alpha, \end{aligned}$$

i.e.

$$a_x = \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M-5m)R}{c^2}} \cos \alpha,$$

and analogously

$$\begin{aligned}
a_y &= \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M-5m)R}{c^2}} \sin \alpha, \\
a_z &= w_x = w_y = 0, \\
w_z &= \frac{i}{c} \frac{\frac{u_y}{ic}}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M-5m)R}{c^2}} \cos \alpha - \frac{i}{c} \frac{\frac{u_x}{ic}}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M-5m)R}{c^2}} \cos \alpha = \\
&= \frac{1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{Gm}{R^2 + \frac{G(M-5m)R}{c^2}} \frac{v_x y - v_y x}{rc^2}.
\end{aligned}$$

In the last equality it is used that $\cos \alpha \approx \frac{x}{r}$, $\sin \alpha \approx \frac{y}{r}$, $u_x \approx v_x \frac{m}{M}$ and $u_y \approx v_y \frac{m}{M}$. We may assume approximately that $x^*/y^* = x/y$, because the barycenter is very close to coordinate origin. Hence $\cos \alpha = x/r$ and $\sin \alpha = y/r$.

By replacing these values in (2.18a,b), and neglecting the terms of order c^{-4} and smaller, we obtain

$$\frac{d^2 x}{dt^2} = \left(1 + \frac{v^2}{c^2}\right) \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M-5m)R}{c^2}} \frac{x - x^*}{R} + \frac{5}{2} \frac{v_x}{c^2} \frac{GM}{R^2} \frac{xv_x + yv_y}{r} - \frac{2v_y}{c^2} \frac{Gm}{R^2} \frac{xv_y - yv_x}{r}, \quad (7.2a)$$

$$\frac{d^2 y}{dt^2} = \left(1 + \frac{v^2}{c^2}\right) \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(M-5m)R}{c^2}} \frac{y - y^*}{R} + \frac{5}{2} \frac{v_y}{c^2} \frac{GM}{R^2} \frac{xv_x + yv_y}{r} + \frac{2v_x}{c^2} \frac{Gm}{R^2} \frac{xv_y - yv_x}{r}, \quad (7.2b)$$

These equations are analogous to (2.2), and continuing to deal like for the perihelion shift in section 2, we obtain

$$\begin{aligned}
\frac{d}{dt} \left(r^2 \frac{d\varphi}{dt} \right) &= \frac{d^2 y}{dt^2} x - \frac{d^2 x}{dt^2} y = \\
&= \frac{5}{2} \frac{GM}{R^2} \frac{xv_y - yv_x}{c^2} \frac{xv_x + yv_y}{r} + 2 \frac{Gm}{R^2} \frac{xv_x + yv_y}{c^2} \frac{xv_y - yv_x}{r} = \\
&= \left(\frac{5}{2} M + 2m \right) \frac{G}{R^2} \frac{xv_x + yv_y}{c^2} \frac{xv_y - yv_x}{r} = - \left(\frac{5}{2} M + 2m \right) \frac{Gr^2}{c^2 R^2} \left(r^2 \frac{d\varphi}{dt} \right) \frac{d}{dt} \left(\frac{1}{r} \right) = \\
&= - \left(\frac{5}{2} M + 2m \right) \frac{GM^2}{c^2 (M+m)^2} \left(r^2 \frac{d\varphi}{dt} \right) \frac{d\rho}{dt},
\end{aligned}$$

where $\rho = 1/r$. The solution of this differential equation is given by

$$r^2 \frac{d\varphi}{dt} = C_2 \left[1 - \left(\frac{5}{2} M + 2m \right) \frac{G}{c^2} \frac{M^2}{(M+m)^2} \rho \right], \quad (7.3)$$

and hence

$$\left(r^2 \frac{d\varphi}{dt} \right)^{-2} \approx C_2^{-2} \left[1 + (5M + 4m) \frac{G}{c^2} \frac{M^2}{(M+m)^2} \rho \right].$$

Analogously to (2.5), now we have

$$\left(\frac{d\rho}{d\varphi} \right)^2 + \rho^2 = v^2 C_2^{-2} \left[1 + (5M + 4m) \frac{G}{c^2} \frac{M^2}{(M+m)^2} \rho \right]. \quad (7.4)$$

Further we should find v as a function of R or r . We can replace

$$\sqrt{1 - \frac{u^2}{c^2}} \approx \left(1 - \frac{v^2}{c^2} \right)^{\frac{u^2}{2v^2}} \approx \left(1 - \frac{v^2}{c^2} \right)^{\frac{m^2}{2M^2}}$$

into the terms of a_x and a_y . According to this approximation, the solution of (2.18d) is given by

$$\begin{aligned} \left(1 - \frac{v^2}{c^2}\right)^{-\frac{1}{2}(1 - \frac{m^2}{M^2})} \left(1 - \frac{m^2}{M^2}\right)^{-1} - \frac{M^2}{(M+m)(M-5m)} \ln\left(1 + \frac{M-5m}{M+m} \frac{GM\rho}{c^2}\right) = \\ = \left(1 - \frac{m^2}{M^2}\right)^{-1} + \frac{Const.}{c^2}. \end{aligned}$$

Hence after some transformations we can express v^2 as a function of ρ up to c^{-2} , and obtain

$$v^2 = 2 \frac{GM^3\rho}{(M+m)^2} - \frac{G^2M^4\rho^2(4M^2 - 4mM - 6m^2)}{c^2(M+m)^4}.$$

Note that the constant of integration *Const.* is neglected in the last formula, because one can verify that it has no role in the final formula for the periastron shift, like for the perihelion shift in section 2. Substituting this value of v^2 into (7.4) we obtain

$$\left(\frac{d\rho}{d\varphi}\right)^2 + \rho^2 = A + B\rho + C\rho^2,$$

where

$$C = (10M + 8m) \frac{G^2M^5}{C_2^2(M+m)^4c^2} - \frac{G^2M^4(4M^2 - 4mM - 6m^2)}{C_2^2(M+m)^4c^2} = \frac{6G^2M^4}{C_2^2(M+m)^2c^2}.$$

Since $C_2 = \frac{2}{P}\pi a^2\sqrt{1-\epsilon^2} = \frac{2\pi}{P}a_r^2\frac{M^2}{(M+m)^2}\sqrt{1-\epsilon^2}$, where a_r is the semi-major axis of the relative orbit and P is the orbit period, we obtain

$$C = \frac{3P^2G^2}{2\pi^2c^2} \frac{(M+m)^2}{a_r^4(1-\epsilon^2)},$$

and analogously to the calculations for the perihelion shift in section 2, for the periastron shift in this case we obtain

$$\Delta\varphi = \frac{3G^2(M+m)^2P^2}{2\pi c^2 a_r^4(1-\epsilon^2)}. \quad (7.5)$$

Using that $a_r^3 = P^2G(M+m)/(4\pi^2)$, we obtain finally

$$\Delta\varphi = \frac{6\pi(4\pi^2)^{1/3}G^{2/3}}{P^{2/3}c^2(1-\epsilon^2)}(M+m)^{2/3}. \quad (7.6)$$

This value (7.6) of the periastron shift is the same which predicts the GR.

Note that if we did not take into account that the pulsar with mass m has an influence on the gravitation from the companion with mass M , i.e. if we omit the expression $1 + \frac{3Gm}{Rc^2}$ in (7.1), then instead of (7.6) the periastron shift would be $\frac{M}{M+m}\Delta\varphi$, where $\Delta\varphi$ is given by (7.6). It means that the companion star participates with $\frac{M}{M+m}\Delta\varphi$ in the total periastron shift given by (7.6). Symmetrically, the pulsar with mass m , participates with $\frac{m}{M+m}\Delta\varphi$ in the total periastron shift.

Now let us assume that

$$a_x = \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(\alpha M + \beta m)R}{c^2}} \frac{x}{r}, \quad \text{and} \quad a_y = \frac{-1}{\sqrt{1 - \frac{u^2}{c^2}}} \frac{GM}{R^2 + \frac{G(\alpha M + \beta m)R}{c^2}} \frac{y}{r}.$$

Then the previous calculation leads that the periastron shift instead of (7.6) is equal to

$$\Delta\varphi = \frac{6\pi(4\pi^2)^{1/3}G^{2/3}}{P^{2/3}c^2(1-\epsilon^2)}(M+m)^{2/3} \times \frac{(7-\alpha)M^2 + (8-\alpha-\beta)mM + (1-\beta)m^2}{6(M+m)^2}.$$

Since the angle $\Delta\varphi$ should be symmetric with respect to M and m , it must be $7 - \alpha = 1 - \beta$, i.e. $\alpha - \beta = 6$. But this is satisfied if and only if the "correcting multiplier" is $1 + \frac{3Gm}{rc^2}$. Further, in that case

$$\Delta\varphi = \frac{6\pi(4\pi^2)^{1/3}G^{2/3}}{P^{2/3}c^2(1-\epsilon^2)}(M+m)^{2/3} \times \frac{7-\alpha}{6},$$

and it depends on the total mass $M+m$, but not on M and m separately. Finally, α must be 1 (and hence $\beta = -5$) in order to be

$$\Delta\varphi = \frac{6\pi(4\pi^2)^{1/3}G^{2/3}}{P^{2/3}c^2(1-\epsilon^2)}M^{2/3}$$

for $m = 0$.

Note that the formula (7.3) does not depend on α and β . Thus, (7.3) for the pulsar's orbit can be important test for ILLTG, like the tests 1 and 2 in section 5. For example, from (7.3) it just follows that

$$\left(r^2 \frac{d\varphi}{dt}\right)_{\text{periastron}} : \left(r^2 \frac{d\varphi}{dt}\right)_{\text{apastron}} = 1 - \frac{(5M+4m)GM^2\epsilon}{a_p c^2 (M+m)^2 (1-\epsilon^2)}, \quad (7.7)$$

where a_p is the semi-major axis of the pulsar's orbit. Using the recent data about the Hulse-Taylor pulsar PSR B1913+16 [12], we get

$$\left(r^2 \frac{d\varphi}{dt}\right)_{\text{periastron}} : \left(r^2 \frac{d\varphi}{dt}\right)_{\text{apastron}} = 1 - 0,47 \times 10^{-5}.$$

Probably using the fact that $\Delta\varphi$ should be a symmetric function with respect to M and m , or some similar conditions, it can be used to obtain the gravitational potential in the form

$$-\frac{GM}{R} + \frac{G^2 M(M-5m)}{2R^2 c^2} - \frac{G^3 M(\alpha M^2 + \beta m M + \gamma m^2)}{3R^3 c^4} + \dots$$

and hence to obtain the post-Newtonian approximations of 2,3,... order. For post-Newtonian approximations of order 2,5; 3,5;... should be considered the multiple moments.

8 ILLTG and the Hubble constant

The representations in sections 1-7 would not be complete if we skip some cosmological consequences considered in [10,11]. In this section the basic arguments of [10] from viewpoint of ILLTG will be considered.

In [10] is given a new view of the Hubble law, showing that the Hubble red shift and the anomalous acceleration of Pioneer spacecraft (see Ref. [1,2,3]) are consequences of the change of the potential in the universe. It is detected red shift such that $\nu = \nu_0 \left(1 - \frac{v}{c}\right)$, and replacing $v = RH$, indeed it is observed that

$$\nu = \nu_0 \left(1 - \frac{RH}{c}\right). \quad (8.1)$$

On the other hand, if the potential changes linearly with the time, after time $t = R/c$ we have additional potential $\frac{R}{c} \frac{\partial V}{\partial t}$, and hence we have a frequency shift such that

$$\nu = \nu_0 \left(1 + \frac{1}{c^2} \frac{R}{c} \frac{\partial V}{\partial t}\right). \quad (8.2)$$

We should remark here that the sign of the potential V is such that near to the spherical gravitational body the potential ($V = \frac{GM}{R}$) is larger than the potential far from gravitation ($V = 0$).

Comparing the formulas (8.1) and (8.2) we obtain

$$H = -\frac{1}{c^2} \frac{\partial V}{\partial t}. \quad (8.3)$$

The change of the potential energy in the universe is probably related with the dark energy, which is about 67% of the whole energy in the universe [6]. Some recent measurements show that H is about 72 km/s/Mpc, but we shall accept value of about 82 km/s/Mpc.

Since $\frac{\partial V}{\partial t} < 0$, in the past the time in the universe was slower than now, and in future it will be faster than now. Indeed, each second the time is faster approximately $\lambda = (1 + H \times 1s) = 1 + \frac{1}{3.76 \times 10^{17}}$ times.

On the other hand, in the last decades are considered the motions of the spacecraft Pioneer 10 and 11, by comparing the initial frequency of a signal which is sent from the Earth to the spacecraft and the frequency of the signal which comes back. The frequency of the received signal on the Earth does not fit with the predicted frequency modeled by using the Doppler effect, the position and motion of the spacecraft using the GR and a lot of perturbations [3]. The observations show that appears almost a constant apparent unmodeled acceleration $a_P \approx (8, 74 \pm 1, 33) \times 10^{-8} \text{cm/s}^2$ which seems to act to the spacecraft toward the Sun.

The acceleration a_P is introduced by formula (15) in [3], i.e.

$$[\nu_{obs}(t) - \nu_{model}(t)]_{DSN} = -\nu_0 \frac{2a_P t}{c}, \quad \text{where } \nu_{model} = \nu_0 \left[1 - 2 \frac{v_{model}(t)}{c} \right], \quad (8.4)$$

where ν_0 is the reference frequency, the factor 2 is because of two way data, v_{model} is the modeled velocity of the spacecraft due to the gravitational and other forces, and ν_{obs} is frequency of the re-transmitted signal observed by DSN antennae. After time $2t$ has been detected a small blue shift on the top of the red shift caused by the motion of the spacecraft outwards the Sun. Form (8.4), an unexplained blue shift

$$\nu = \nu_0 \left(1 + \frac{2a_P t}{c} \right), \quad (8.5)$$

is detected, where $2t$ is the time of the light signal in two directions. Without using the acceleration a_P , the anomalous frequency shift can be interpreted by "clock acceleration" $-a_t = -2, 8 \times 10^{-18} \text{s/s}^2$ [3]. This causes just the blue shift given by (8.5). This model assumes that the time is uniform, but there is only technical problem with the clocks. Although this model gives good explanations for both frequency shift and the trajectories, it is rejected [3].

Many people have noticed that the acceleration a_P and the Hubble constant H are related by [3]

$$a_P = cH. \quad (8.6)$$

Indeed, if we assume that $H=82 \text{km/s/Mpc}$, then from (8.6) it follows $a_P = 8 \times 10^{-8} \text{cm/s}^2$. Thus we assumed that $H=82 \text{km/s/Mpc}$.

Now we shall explain the apparent acceleration a_P connected with the frequency shift, according to ILLTG. We denote by x, y, z, t the orthonormal coordinates and denote by X, Y, Z, T ordinary coordinates where there is time dependent gravitational potential $V = -c^2 H t$. According to (1.1), where $V = -H c^2 t$, we have the following equalities

$$\begin{aligned} dx &= \left(1 + \frac{V}{c^2} \right)^{-1} dX = (1 + tH) dX, & dy &= \left(1 + \frac{V}{c^2} \right)^{-1} dY = (1 + tH) dY, \\ dz &= \left(1 + \frac{V}{c^2} \right)^{-1} dZ = (1 + tH) dZ, & dt &= \left(1 - \frac{V}{c^2} \right)^{-1} dT = (1 - tH) dT. \end{aligned} \quad (8.7)$$

Remark. Note that it is sufficient in this section to assume that (8.7) are satisfied. Then it is not necessary to speak about the time dependent gravitational potential. The coefficients $1 + Ht$ and $1 - Ht$ in (8.7) should be exponential functions, but neglecting H^2 and smaller quantities we accept these linear functions.

If we use orthonormal coordinates x, y, z, t , then the motion of the spacecraft does not depend on the time dependent potential $-H c^2 t$. Indeed, it is easy to see that the tensor F_{ij} does not depend on this potential, and hence the motion of the particles (2.2) does not depend on the time dependent

potential, which means that $(\frac{d^2x}{dt^2}, \frac{d^2y}{dt^2}, \frac{d^2z}{dt^2}) = (0, 0, 0)$, if we consider only the time dependent potential. Using the equalities (8.7), (2.2') and (8.6) we have an additional 3-acceleration

$$-3HT\left(\frac{d^2X}{dT^2}, \frac{d^2Y}{dT^2}, \frac{d^2Z}{dT^2}\right) - 2\left(H\frac{dX}{dT}, H\frac{dY}{dT}, H\frac{dZ}{dT}\right). \quad (8.8)$$

In the explanation of the shift frequency of Pioneers, the first component $-3HT\left(\frac{d^2X}{dT^2}, \frac{d^2Y}{dT^2}, \frac{d^2Z}{dT^2}\right)$ is smaller, compared with the second and main component

$$-2\left(H\frac{dX}{dT}, H\frac{dY}{dT}, H\frac{dZ}{dT}\right) = -2\left(\frac{a_P}{c}\frac{dX}{dT}, \frac{a_P}{c}\frac{dY}{dT}, \frac{a_P}{c}\frac{dZ}{dT}\right).$$

We shall explain why it is measured the blue shift given by (8.5), instead of the red shift like in (8.1). It will be explained without using the acceleration (8.8), but only the definition (8.4) where a_P is introduced, and our model of time dependent gravitational potential.

Let $R = R(T)$ be the distance from the spacecraft to the DSN antenna. Let us calculate $\frac{\Delta\nu}{\nu_0} = (\nu_{obs} - \nu_0)/\nu_0$. According to our model, neglecting the terms containing H^2 , this expression is equal to

$$\frac{\Delta\nu}{\nu_0} = -2H\frac{R}{c} - 2\frac{dR}{dT},$$

where the expression $-2H\frac{R}{c}$ corresponds to the Hubble red shift from (8.2). On the other side, according to (8.4), where v_{model} it determined without using the Hubble constant H , we have

$$\frac{\Delta\nu}{\nu_0} = 2a_P\frac{R}{c^2} - 2\left(\frac{dR}{dT}\right)_{H=0}.$$

From the last two equalities we are able to find the expression for a_P

$$a_P = -c\left(\frac{1}{R}\frac{dR}{dT} - \frac{1}{R}\left(\frac{dR}{dT}\right)_{H=0} + H\right).$$

Using that $\frac{dR}{dT} = \frac{dr}{dt}(1 - 2HT) = \left(\frac{dR}{dT}\right)_{H=0}(1 - 2HT)$ according to (8.7), we obtain finally

$$a_P = cH\left(2\frac{dR}{dT}\frac{T}{R} - 1\right). \quad (8.9)$$

The equality (8.9) gives the required expression of the acceleration a_P . Now we are not able to determine the initial value of T , i.e. when we should start to measure the time T in (8.9). It depends on the estimation of the initial values for the motion of the spacecraft, i.e. it depends on the departure of the real initial values. Thus, for different spacecraft are obtained close but different values of a_P . Note also that for each initial value of T , when T tends to infinity (or R tends to infinity), then a_P tends to cH . In an ideal case, if the experiment is done in an inertial system, then $dR/dT = R/T$ is a constant velocity and then $a_P \equiv cH$.

The change of the gravitational potential causes slight changes in the results of the known experiments about GR, but they are negligible. The change of the results in the test of Shapiro time delay is negligible if we consider radio signals on short distances like in the solar system. Indeed, the change of the potential in the universe is about $2,4 \times 10^3 \text{cm}^2/\text{s}^2$ each second, while the gravitational potential on 1AU from the Sun is equal to $30^2 \text{km}^2/\text{s}^2 = 9 \times 10^{12} \text{cm}^2/\text{s}^2$. This potential difference will be achieved from the universe after 118 years, which is too long compared with the time needed for an experiment about Shapiro time delay. The change of the deflection of the light rays near the Sun is completely negligible up to c^{-2} . The angle between two perihelions is changed additionally of order $\frac{\Theta^2 H^2}{\epsilon}$, where Θ is the orbit period of the planet and ϵ is the eccentricity of the elliptic orbit. This change of the angle is negligible with respect to the measured angle $\frac{6GM\pi}{ac^2(1-\epsilon^2)}$.

Note that if we replace V by $V - c^2 H t$ ($tH \ll 1$) in the metric of GR

$$(g_{ij}) = \text{diag}\left(-1 + \frac{2V}{c^2} - \frac{2V^2}{c^4}, 1 + \frac{2V}{c^2}, 1 + \frac{2V}{c^2}, 1 + \frac{2V}{c^2}\right)$$

the additional acceleration is not convenient. The situation becomes much worse if t is very large. While the Newton's theory is invariant under the gauge transformation $V \mapsto V + C$, and ILLTG is invariant under the gauge transformation $\mu \mapsto C\mu$, GR does not possess similar property. The equations of motion are unchanged if we replace the metric g_{ij} by Cg_{ij} , but it can not be correlated with the change of the gravitational potential.

References

- [1] J. D. Anderson, S. Turyshev and M. M. Nieto, Effect of the Pioneer Anomaly on Long-Period Comet Orbits, AAS 201st Meeting, January, 2003, *Bulletin of the American Astronomical Society* **34**, # 4.
- [2] J.D. Anderson, P. A. Laing, E. L. Lau, A. S. Liu, M. M. Nieto and S. G. Turyshev, Indication, from Pioneer 10/11, Galileo and Ulysses Data, of an Apparent Anomalous, Weak, Long-Range Acceleration, *Phys. Rev. Lett.* **81**, 2858 (1998).
- [3] J. D. Anderson, P. A. Laing, E. L. Lau, A. S. Liu, M. M. Nieto and S. G. Turyshev, Study of anomalous acceleration of Pioneer 10 and 11, *Phys. Rev. D* **65**, 082004.
- [4] B. Bertotti, L. Iess and P. Tortora, A test of general relativity using radio links with the Cassini spacecraft, *Nature* **425**, 374-376.
- [5] D. F. Lawden, *Tensor Calculus and Relativity*, Chapman and Hall, London, 1976.
- [6] W. L. Freedman and M. S. Turner, Measuring and understanding of the universe, *Rev. of Modern Phys.* **75**, 1433-1447 (2003).
- [7] J. M. Overduin and P. S. Wesson, *Phys. Rept.*, **283**, 303 (1997).
- [8] K. Trenčevski, One model of gravitation and mechanics, *Tensor N.S.* **53**, 70-82 (1993).
- [9] K. Trenčevski and V. Balan, On the shrinking of rotational configurations and the corresponding inertial forces, submitted for publication.
- [10] K. Trenčevski, Hubble red shift and the anomalous acceleration of Pioneer 10 and 11, submitted for publication, eprint gr-qc/0402024.
- [11] K. Trenčevski, Deformation of the planetary orbits caused by the time dependent gravitational potential in the universe, submitted for publication, eprint gr-qc/0403067.
- [12] J. M. Weisberg, J. H. Taylor, The relativistic Binary Pulsar B1913+16, 2003, in ASP Conf. Ser. 302, p. 93., eprint: astro-ph/0211217.